




documenta

naturae

No. 87

MÜNCHEN 1994

PALYNOLOGICAL ASPECTS OF THE PERMO - TRIASSIC SUCCESSION IN THE HOLY CROSS MOUNTAINS, POLAND



DOCUMENTA NATURAE

Nr. 87

1994

ISSN

0723-8428

Herausgeber:

Dr. Hans-Joachim Gregor, Palsweiserstr. 5m, D-82140 Olching
Dr. Heinz J. Unger, Nußbaumstraße 13, D-85435 Altenerding

Die Zeitschriftenreihe erscheint in zwangloser Folge mit Themen aus den Gebieten Geologie, Paläontologie, Botanik, Anthropologie, Vor- und Frühgeschichte, Domestikationsforschung, Stratigraphie, Lagerstättenkunde usw.

Die Zeitschriftenreihe ist auch Mitteilungsorgan der Paläobotanisch-Biostratigraphischen Arbeitsgruppe (PBA).

Für die einzelnen Beiträge zeichnen die Autoren verantwortlich, für die Gesamtgestaltung die Herausgeber.

Überweisung der Schutzgebühr erbeten auf das Konto 6410317280 bei der Bayerischen Hypotheken- und Wechselbank München (BLZ 700 200 01) - Inh. H.-J. Gregor

Bestellungen bei Buchhandlungen und den Herausgebern.

Copyright: bei Verlag und Autoren

Umschlagbild: H.-J. Gregor und Autor(en)

documenta naturae

87

1-76

4 Abb.

3. Tab.

7. Taf.

München 1994

**Palynological aspects of the Permo-Triassic succession
in the Holy Cross Mountains, Poland**

with 4 Figures, 3 Tables and 7 Plates

Anna Fijałkowska*

Address of the author

*State Geological Institute, Zgoda 21, 25-395 Kielce, Poland

Summary

A palynostratigraphic study of the Permo-Triassic sequence in the Holy Cross Mountains resulted in the identification of four spore-pollen assemblages which represent *Lueckisporites virkkiae* and LT1 palynological zones.

The correlation was drawn between the Holy Cross spore-pollen assemblages and the contemporaneous spectra known from the other European regions assigned both to the German and Alpine basins.

The paleoclimatic model obtained from this study indicates the consequent aridation during the Late Permian, documented by the distinct domination of the xerophytic elements in the miosores spectra and an increase of humidity in the Early Triassic.

Streszczenie

W wyniku badań palinologicznych osadów zechsztynu na obszarze Górz Świętokrzyskich wyróżniono trzy zespoły sporowo-pyłkowe należące do późnopermskiej zony *Lueckisporites virkkiae* oraz jeden zespół reprezentujący wczesnotriasową zonę LT1.

Przeprowadzono korelację między zespołami świętokrzyskimi a równowiekowymi spekrami znanyimi z innych obszarów Europy, należących zarówno do basenu germanickiego jak i alpejskiego.

Paleoklimatyczny model otrzymany w wyniku tych badań wskazuje na panowanie suchego klimatu w czasie późnego permu, udokumentowanego wyraźną dominacją kserofitycznych elementów w obrębie zespołów miospor oraz na wzrost wilgotności we wczesnym triasie.

Table of contents

Summary	3
Introduction	4
Methods and materials	5
General characteristic of the Zechstein and Lower Buntsandstein lithological complexes	6
Systematics	10
Description of the new species, new forms and some new combinations	18
Palynostratigraphy	22
Relation between microfloristic assemblages and lithostratigraphy	34
Boundary problem	36
Paleoclimate and paleoenvironment	37
Conclusions	40
Acknowledgements	41
References	42
Appendix 1a-d. Range charts	52
Appendix 2. Alphabetical list of all identified miospore taxa	58
Appendix 3. Taxonomy of selected taxa	60
Explanation to the Plates	68

INTRODUCTION

This paper presents the first completely, as far as the geological conditions allowed, palynozonation for the Permo-Triassic sequence in the Holy Cross Mountains. This zonation is applied to the boundary problem.

The Upper Permian (Zechstein) - Lower Triassic (Lower Buntsandstein) section in the Holy Cross Mts has been known from several boreholes (Fig. 1). All palynological data come from the borehole samples.

The currently used lithological scheme of the Polish Zechstein, based on Richter-Bernburg's (1955) sedimentological cycles, was established by Wagner, Piątkowski and Peryt (1978) and adopted to the Holy Cross Mountains by Kowalczewski (1978), Pawłowska (1978), Rup (1985) and Kowalczewski and Rup (1989).

Kuleta (1985, 1990) proposed the lithostratigraphic framework for the Lower Buntsandstein in this area.

Any faunistic studies, with an exception of foraminifera from the Lower Zechstein elaborated by Jurkiewicz (1962, 1966), didn't bring the satisfactory results. Therefore the palynological research seemed to be the only chance for the creation of biostratigraphical scheme. It was begun by Kotańska and Krason (1966) who listed some species of the Upper Permian miospores from Gałżice region (SW part of the Holy Cross Mts). These studies were continued by Dybova-Jachowicz and Laszko (see Dybova-Jachowicz, 1976; Dybova-Jachowicz, Laszko, 1976; Dybova-Jachowicz, Jagielska, 1977; Dybova-Jachowicz, Laszko, 1978, 1980; Dybova-Jachowicz, 1981) who finally distinguished one spore-pollen assemblage in the Zechstein and two - in the Middle and Upper Buntsandstein. The palynological documentation of the Zechstein-Buntsandstein boundary has been an open problem till the author's investigations. They indicated palynological hiatus at the boundary which was considered to be lithological continuity (Kuleta, 1985; Pieńkowski, 1989).

The Zechstein-Lower Buntsandstein palynological assemblages have been for the first time used as paleoenvironmental and paleoclimatological indicators. Their interpretation should help to understanding the geological events in the Holy Cross Mountains.




Fig. 1.
Location map of boreholes

METHODS AND MATERIALS

Palynological data from 22 boreholes was compiled in this study (Fig. 1). The total 680 samples were taken from the selected sections at intervals about 200 m. The summary length of the studied core was 4050 m. Finally the 285 samples appeared positive.

Samples were collected mainly from the black, dark grey and greenish mudstones, siltstones, fine grained clayey sandstones, limestones and marls. Almost all samples from the detritic red, reddish and mottled deposits were negative.

Samples of material weighing 30-50 g were crumbled mechanically and then treated with the following reagents: 1) 10% HCl; 2) 40% HF in the cold state for a period of 3-7 days; 3) washed with a distilled water and treated at 70 C for 2 hours in 10% HCl; 4) floated in heavy liquid CdJ₂ + KJ₂, with a specific weight of 2,1.

The organic material thus obtained was oxidated in 30% HNO for 24 hours. The residuum was moluted in glycerine gelly.

The quantity palynological analysis was based on a count of all specimens recognized in slide which were fixed as 100%. Microplankton was counted separately.

GENERAL CHARACTERISTIC OF THE ZECHSTEIN AND LOWER BUNTSANDSTEIN LITHOLOGICAL COMPLEXES

The Holy Cross Mts belonged to the coastal zone of the German basin during the Late Permian and Early Triassic. Therefore some members of the Zechstein and Buntsandstein lithological sections are missed or reduced in comparison to the deeper part of basin.

The following lithological complexes are currently distinguished within the Zechstein on the so-called "near" Permian-Mesozoic edge of the Holy Cross Paleozoic core:

Zechstein Limestone (Ca1) locally underlaying by the Copper-Bearing Shale (T1), Lower Anhydrite (A1d) in some sections, Terrigenous Series (T1r) and Upper Anhydrite (A1g) which all form the Werra Cycle (PZ1); the Strassfurt Cycle (PZ2) is developed as the Terrigenous Series T2r; the Grey Pelite (T3), Platy Dolomite (Ca3) and only locally Main Anhydrite (A3) represent the Leine Cycle (PZ3). The Uppermost Zechstein deposits consist of the Top Terrigenous Series (Pzt) (Tab.1).

In those Zechstein sections, in which both the sulphates of PZ1 Cycle and the chemical deposits of the PZ3 Cycle are missing, all the rocks overlaying the Zechstein Limestone have been defined as Terrigenous Upper Permian Series (P2t).

The boundary between Zechstein and Buntsandstein has been determinated within the "variegated" clastic series only on the basis of lithological criteria, and thus should be regarded as hypothetical (Kowalczewski and Rup, 1989).

Three complexes were defined in the Lower Buntsandstein deposits: A0 ("knobby"), A1 and B (Kuleta, 1985) (Tab. 1).

The PZ1 deposits are connected with the maximum spreading of the Zechstein basin at the studied area (Wagner et al., 1978) and run into the maximum thickness of 54,0-179,5 m among the Upper Permian rocks (Rup, 1985). They usually begins with the Zechstein Limestone (Ca1). It is only in few boreholes that could be distinguished thin layers

(1,5-2,5 m) corresponding to the horizon of the Copper-Bearing Shale (T1). These are marly dolomites and argillaceous mudstones with sulphide mineralization. In the other sections the Zechstein Limestone lies on the differentiated, Paleozoic basement of the Cambrian, Silurian, Devonian and Lower Carboniferous rocks. It is very often underlined by the conglomerates

		B	varigrained sandstones and mudstones building fining upward cycles
	Tp1	A1	sandstones (locally with conglomerates at the base) alternated with mudstones
		A0	mudstones with the carbonate nodules
	Pzt	Top Terrigenous Series Pzt	sandstones and mudstones with the conglomerates intercalations
			mudstones
	Pz3	Main Anhydrite A3	anhydrites alternated with mudstones
		Platy Dolomite Ca 3	dolomites and laminated limestones
		Grey Pelite T3	sandstones, mudstones and pelites
E - I	Pz2	Terrigenous Series T2r	mudstones and sandy mudstones
S - T	T	Upper Anhydrite Alg	anhydrites and gypsum alternated with mudstones
C - H		Terrigenous Series T1r	sandy mudstones (locally with the carbonate nodules) with the pseudomorphs after sulphates
Z - E	Pz1	Lower Anhydrite Ald	anhydrites alternated with mudstones
		Zechstein Limestone Ca 1	limestones and oolithic dolomites silty limestones or dolomites
		Copper Shale T1	mudstones

Tab. 1.

Lithostratigraphy of the Zechstein and Lower Buntsandstein in the Holy Cross Mountains.

of differentiated lithology and thickness. Besides the most common variant composed of the pebbles of the Devonian limestones and dolomites there occur conglomerates built of the Cambrian and Lower Devonian sandstones. The thickness of these conglomerates varies from few to 200 m. The problem of their stratigraphical position has not been solved so far. According to some authors (Rup, 1985) they belong to the Lower Permian but their top part probably represent so-call "basement conglomerate" of the Upper Permian age (Pawlowska, 1964; Czarniecki et al., 1965).

The Zechstein Limestone is an extensive complex of carbonate rocks, up to 30-100 m in thickness, consisting of micritic, oolitic limestones and dolomites and marls. More detritic material is observed in the sections located near the Paleozoic base. The microfacies show that this complex was formed in a lagoon zone and in a barrier developed at the edge of the carbonate shelf platform (Kowalczewski and Rup, 1989).

The oldest sulphate rocks of the PZ1 Cycle, correlated with the Lower Anhydrite (A1d) were found over Ca1 rocks only in five boreholes. These are anhydrites and gypsum alternated with siltstones, mudstones or muddy dolomites of thickness 3-6 m.

Above these anhydrites, and - in the other sections - above Ca1 deposits, there are mainly detrital rocks, defined as recessive Terrigenous Series (T1r), up to 7,5-42,2 m thick. These are calcareous and dolomitic mudstones with intercalations of fine sandstones and - less frequently - of claystones, marls and dolomitic marls in the northern subregion. In the southern subregion occur mainly brown mudstones with carbonate nodules. There is a sedimentary breccia in some boreholes composed of fragments of marly and dolomitic limestones and mudstones, embedded in an argillaceous matrix. In the southern part of the studied area, this breccia is replaced by the intercalations of fine conglomerates (Kowalczewski, Rup, 1989).

In the top parts of the PZ1 cycle there are sulphate rocks which correspond to the Upper Anhydrite (A1g). The thickness of this series varies from 2 to 28 m. In the majority of boreholes this series is bipartite, divided by the thin, several meters thick layer of brown claystones or mudstones. Its lower part is composed of anhydrites laminated with clay, the upper one is distinctly dominated by sulphates. Equivalents of these rocks are mudstones with the thin layers of algal limestones in the southern subregion. In the A1g deposits there is also a sedimentary breccia corresponding most probably to the Upper Anhydrite Breccia (BrA1g).

The supposed deposits of the Strassfurt Cycle (PZ2) are represented only by detrital

rocks - sandstones and mudstones referred to the recessive Terrigenous Serie (T2r) of thickness 4,5-27,5 m.

The Leine Cycle (PZ3) sediments occur in most of the sections and they are better developed than those of the PZ2 Cycle. Rocks, that are equivalent to the Grey Pelite (T3), can be distinguished in some sections. These are grey, strongly calcareous mudstones and siltstones 0,6-7,5 m thick. In the W part of the investigated area the base of this series is composed of grey, fine sandstones and sandy mudstones which becomes more conglomeratic closed to the Paleozoic base. It creates an important correlation horizont (Kowalczewski, Rup, 1989; Rup, 1985).

The equivalent of the Platy Dolomite (Ca3) is more widely distributed. In the southern subregion its thickness does not exceed 2m; it occurs there in the form of the algal limestones with an admixture of bioclasts. These deposits probably originated in the intertidal zone (Rup, 1985). In the northern subregion Ca3 deposits are thicker (up to 10 m) and bipartite: algal biolitites at the bottom are overlain by the alternating micrite and marl layers.

The equivalent of the Main Anhydrite (A3) was distinguished only in the four boreholes. These are sulphates, mainly anhydrites, laminated with dolomicrites and dolosparites of the total thickness of 1,0-3,1 m (Rup, 1985).

All the rocks above Ca3 and A3 deposits are regarded as the Top Terrigenous Series (Pzt). They are composed of the brown sandstones and sandy mudstones with the carbonate nodules and - less often - of claystones and conglomerates near the Paleozoic core. Their thickness varies from 10,9 to 53,7 m (Kowalczewski, Rup, 1989; Rup, 1985).

The accumulation of the Permian deposits in the area of the Holy Cross Mts was closely connected with the relief of the terrain in and tectonic activity of that region during the late Variscan movements. The Cambrian-Lower Carboniferous basement which had been folded during the Upper Carboniferous was deeply graded before the Zechstein. The relief of the terrain reached its mature stage at that time. Towards the end of the Zechstein the exposed Paleozoic core which had not been covered by the Zechstein transgression reached its senile stage.

During the Permian the Holy Cross Mts were subject to slow elevating movements. It was only three times that these movements were stopped by the short-term lowering trends. These lowering movements led to marine transgression of the Cycles PZ1 and PZ3 which reached the "near" margin of the Holy Cross Mts and to the transgression of Cycle PZ2

which covered the "distant" peripheries of this region. The lowering pulsations were marked by varying intensity and they lasted on this area for a longer (PZ1) and shorter (PZ3) period (Kowalczewski, Rup, 1989).

The following lithological complexes are distinguished within the Lower Buntsandstein: A0, A1 and B. The lowermost A0 complex lies in the sedimentological continuity with the Zechstein deposits. It is composed of brown, structureless mudstones and sandy mudstones containing the carbonate nodules and irregular concentriations. The thickness of this series varies from 9,6 to 38, 0 m (Kuleta, 1985). This complex originated, according to Pieńkowski (1989), in the playa like environment and was an echo of the Lower Triassic transgression in the Central Poland.

The A1 complex was identified in the seven boreholes. It is formed by the reddish and brown, sandy and muddy deposits (20,5-67,0 m thick) with the conglomerates and coarse sandstone at the bottom. In the top there dominate siltstones and mudstones with the wavy sedimentary structures which show the marine character of the deposit (Kuleta, 1985).

The B complex is the main member of the Lower Buntsandstein section in this area. It reaches the maximum thickness of 68,0- 167,0 m. Lithologically is monotonous, built of pink-reddish, varie -grained, sometimes conglomeratic sandstones. The cyclity of sedimentation with the upward fine-grained sequences is the most characteristic feature of this deposit. The other sedimentological structures as large-scale diagonal bedding indicate the river environment of the deposition (Kuleta, 1985).

SYSTEMATICS

In total 90 species of miospores belonged to 34 genera and 7 species of acritarchs were identified in the investigated material. Miospores were classified according to the systematics presented by Potonié (1956, 1958, 1960, 1970) and Potonié and Kremp (1954, 1955, 1956a, 1956b) completed with the following papers: Dybova and Jachowicz (1957), Jansonius (1962), Klaus (1963) Hart (1964, 1965) Visscher (1966), Scheuring (1970), Dibner (1971) and Foster (1983).

A new form *Laevigatisporites giganteus* Dybova et Jachowicz f. *microsignus* was distinguished within the spores and a new combination - *Lycospora permica* (Inosova, 1976) was proposed. A new species *Illinites kosankei* was described among the bisaccate pollen

grains and few new combinations were introduced into *Lunatisporites* (Leschik, 1956) Scheuring 1970 genus: *L. alatus* (Klaus, 1963), *L. gracilis* (Jansonius, 1962), *L. labdacus* (Klaus, 1963), *L. obex* (Balme, 1963) and *L. transversundatus* (Jansonius, 1962). A new species name *Lunatisporites microsaccatus* was proposed for the form *Taeniaesporites* sp. U Jansonius 1962 and *Triadispora visscheri* - for *Triadispora muelleri* Visscher, 1966.

LIST OF THE DETERMINED SPECIES

Anteturma Proximegerminantes Potonié, 1970

Turma Triletes-Azonales Potonié, 1970

Subturma Azonotriletes Luber, 1935

Infraturma Laevigati (Bennie et Kidston, 1886) Potonié, 1956

Genus CALAMOSPORA Schopf, Wilson et Bentall, 1944

Calamospora pedata Kosanke, 1950 - Pl. I, fig. 2

Calamospora plicata (Luber et Waltz, 1941) Hart, 1965 - Pl. I, fig. 1

Genus LAEVIGATISPORITES Dybova et Jachowicz, 1957

Laevigatisporites giganteus Dybova et Jachowicz, 1957 f. *microsignus* f.nov. - Pl. I, fig. 5

Laevigatisporites minimalis f. *pulla* Dybova et Jachowicz, 1957 - Pl. I, fig. 4

Genus PUNCTATISPORITES (Ibrahim, 1933) Potonié et Kremp, 1954

Punctatisporites triassicus Schulz, 1964 - Pl. I, fig. 8

Infraturma Apiculati (Bennie et Kidston, 1886) Potonié, 1958

Genus APICULATISPORITES (Ibrahim, 1933) Potonié et Kremp, 1956

Apiculatisporites apiculatus f. *media* Dybova et Jachowicz, 1957 - Pl. I, fig. 9

Genus CYCLOTRILETES Mädler, 1964

Cyclogtriletes microgranifer Mädler, 1964 - Pl. I, fig. 3

Cyclogtriletes oligogranifer Mädler, 1964 - Pl. I, fig. 7

Infraturma Verrucati Dybova et Jachowicz, 1957

Genus VERRUCOSISPORITES (Ibrahim, 1933) Potonié et Kremp, 1956

Verrucosporites pseudomorulae Visscher, 1966 - Pl. I, fig. 6

Rodzaj **GUTTATISPORITES** Visscher, 1966

Guttatisporites sp. - Pl. I, fig. 15

Turma Zonales (Bennie et Kidston, 1886) Potonié, 1956

Subturma Zonotriletes (Waltz, 1935) Potonié et Kremp, 1954

Infraturma Cingulati Potonié et Kremp, 1954

Genus LYCOSPORA Schopf, Wilson et Bentall, 1944

Lycospora permica (Inosova, 1976) comb. nov. - Pl. I, fig. 12

Genus DENSOISPORITES (Weyland et Krieger, 1953) Dettmann, 1963

Densoisporites playfordii (Balme, 1963) Dettmann, 1963 - Pl. I, fig. 17

Genus LUNDBLADISPORA (Balme, 1963) Playford, 1965

Lundbladispora brevicula Balme, 1963 - Pl. I, fig. 13

Lundbladispora cf. *obsoleta* Balme, 1970 - Pl. I, fig. 14

Turma Monoletes Ibrahim, 1933

Subturma Azonomonoletes Luber, 1935

Infraturma Laevigatimonoletes Dybova et Jachowicz, 1957

Genus LAEVIGATOSPORITES Ibrahim, 1933

Laevigatosporites vulgaris (Ibrahim, 1932) Ibrahim, 1933 f. *minor* Loose, 1934 - Pl. I, fig. 11

Anteturma Variegerminantes Potonié, 1970

Turma Saccites Erdtman, 1947

Subturma Monosaccites (Chitaley, 1951) Potonié et Kremp, 1954

Infraturma Monopolsacciti (Hart, 1965) Dibner, 1970

Subinfraturma Proximalsaccini Dibner, 1970

Genus FLORINITES Schopf, Wilson et Bentall, 1944

Florinites sp. - Pl. I, fig. 10

Genus *PERISACCUS* (Naumova, 1953) Potonié, 1958 emend. Klaus, 1963

Perisaccus granulatus Klaus, 1963 - Pl. I, fig. 16

Subinfraturma **Distalsaccini** Dibner, 1970

Genus *ENDOSPORITES* Wilson et Coe, 1940

Endosporites hexarecticulatus Klaus, 1963 - Pl. I, fig. 19

Genus *POTONIEISPORITES* Bharadwaj, 1954

Potonieisporites simplex Wilson, 1962 - Pl. I, fig. 18

Infraturma **Dipolsacciti** (Hart, 1965) Dibner, 1970

Subinfraturma **Parasaccini** (Maheshwari, 1967) Dibner, 1970

Genus *CORDAITINA* (Samoilovich, 1953) Hart, 1963

Cordaitina donetziana Inosova, 1976 - Pl. I, fig. 19, 20

Cordaitina uralensis (Luber, 1941) Dibner, 1970 - Pl. II, fig. 7

Genus *NUSKOISPORITES* Potonié et Klaus, 1954

Nuskoisporites dulhuntyi Potonié et Klaus, 1954 - Pl. II, fig. 4

Nuskoisporites klausii Grebe, 1957 - Pl. II, fig. 5

Subinfraturma **Apertacorpini** Dibner, 1970

Genus *TRIZONAESPORITES* (Leschik, 1956) Klaus, 1963

Trizonaesporites grandis Leschik, 1956 - Pl. II, fig. 1,3

Genus *PLICATIOPOLLENITES* Lele, 1964

Plicatipollenites indicus Lele, 1964 - Pl. II, fig. 2

Subinfraturma **Amphisaccini** (Lele, 1965) Dibner, 1970

Genus *CRUCISACCITES* Lele et Maithy, 1964

Crucisaccites cf. latisulcatus Lele et Maithy, 1964 - Pl. II, fig. 6

Subturma Disaccites Cookson, 1947

Infraturma Striatiti Pant, 1954

Genus **PROTOHAPLOXYPINUS** (Samoilovich, 1953) Hart, 1964 emend. Morbey, 1975

Protohaploxyypinus jacobii (Jansonius, 1962) Hart, 1964 - Pl. III, fig. 3

Protohaploxyypinus cf. latissimus (Luber et Waltz, 1941) Samoilovich, 1953 - Pl. III, fig. 2

Protohaploxyypinus pantii (Jansonius, 1962) Orłowska-Zwolińska, 1984 - Pl. III, fig. 6

Protohaploxyypinus cf. rhomboeformis (Polukhina, 1960) Hart, 1964 - Pl. III, fig. 4

Protohaploxyypinus samoilovichii (Jansonius, 1962) Hart, 1964 - Pl. III, fig. 1

Genus **STROTHERSPORITES** (Wilson, 1962) Klaus, 1963

Strotersporites richteri (Klaus, 1955) Wilson, 1962 - Pl. III, fig. 8

Strotersporites wilsoni Klaus, 1963 - Pl. III, fig. 11

Genus **STRIATOPODOCARPITES** (Zaricheva et Sedova, 1956) Hart, 1964

Striatopodocarpites sp. - Pl. III, fig. 13

Genus **STRIATOABIETITES** (Sedova, 1956) Hart, 1964

Striatoabietites aytugii (Visscher, 1966) Scheuring, 1970 - Pl. III, fig. 5

Striatoabietites balmei Klaus, 1964 - Pl. III, fig. 7

Genus **LUECKISPORITES** (Potonié et Klaus, 1954) Jansonius, 1962

Lueckisporites virkkiae Potonié et Klaus, 1954 - Pl. VII, fig. 1-6, 8-9

Lueckisporites virkkiae Potonié et Klaus, 1954 with a "dark body" f. nov. - Pl. VII, fig. 7

Genus **LUNATISPORITES** (Leschik, 1955) Scheuring, 1970

Lunatisporites acutus Leshik, 1955 - Pl. III, fig. 14

Lunatisporites alatus (Klaus, 1963) comb. nov. - Pl. III, fig. 9

Lunatisporites gracilis (Jansonius, 1962) comb. nov. - Pl. III, fig. 10

Lunatisporites hexagonalis (Jansonius, 1962) Scheuring, 1970 - Pl. III, fig. 15

Lunatisporites labdacus (Klaus, 1963) comb. nov. - Pl. III, fig. 16

Lunatisporites microsaccatus (Jansonius, 1962) comb. nov. Pl. III, fig. 17

Lunatisporites multiplex (Visscher, 1966) Scheuring, 1970 - Pl. IV, fig. 7

Lunatisporites noviaulensis (Leschik, 1956) Scheuring, 1970 - Pl. IV, fig. 1

Lunatisporites obex (Balme, 1963) comb. nov. - Pl. III, fig. 12

Lunatisporites ortisei (Klaus, 1963) Góczán, 1987 - Pl. III, fig. 18

Lunatisporites transversundatus (Jansonius, 1962) comb. nov. - Pl. IV, fig. 4

Genus *PROTOSACCULINA* (Maliavkina, 1953) Jansonius, 1962

Protosacculina sp. - Pl. IV, fig. 14

Genus *VITTATINA* (Luber, 1941) Wilson, 1962

Vittatina costabilis Wilson, 1962 - Pl. IV, fig. 3

Vittatina hiltonensis Chaloner et Clarke, 1962 - Pl. IV, fig. 5

Vittatina subsaccata Samoilovich, 1953 - Pl. IV, fig. 2, 6

Vittatina vittifera (Luber et Waltz, 1941) Samoilovich, 1953 - Pl. IV, fig. 11

Genus *HAMIAPOLLENITES* Wilson ex Jansonius, 1962

Hamiapollenites cf. *bifurcatus* Jansonius, 1962 - Pl. IV, fig. 12

Infraturma **Disacciatrileti** Leschik, 1956

Genus *KLAUSIPOLLENITES* Jansonius, 1962

Klausipollenites decipiens Jansonius, 1962 - Pl. IV, fig. 15

Klausipollenites minimus Góczán, 1987 - Pl. IV, fig. 9-10

Klausipollenites schaubergeri (Potonié et Klaus, 1954) Jansonius, 1962 - Pl. IV, fig. 8

Klausipollenites staplinii Jansonius, 1962 - Pl. IV, fig. 13

Genus *FALCISPORITES* (Leschik, 1956) Klaus, 1963

Falcisporites snopkove Visscher, 1966 - Pl. IV, fig. 11

Falcisporites zapfei (Potonié et Klaus, 1954) Leschik, 1955 - Pl. IV, fig. 16

Genus *VESICASPORA* Schemel, 1951

Vesicaspora schemeli Klaus, 1963 - Pl. IV, fig. 17

Genus *PARAVESICASPORA* Klaus, 1963

Paravesicaspora splendens Klaus, 1963 - Pl. IV, fig. 20

Genus *PLATYSACCUS* (Naumova, 1937) Potonié et Klaus, 1954

Platysaccus leschiki Hart, 1960 - Pl. V, fig. 1

Platysaccus niger Mädler, 1964 - Pl. V, fig. 3

Platysaccus papilionis Potonié et Klaus, 1954 - Pl. V, fig. 2

Infraturma *Disaccitrileti* Leschik, 1956

Genus *ILLINITES* (Kosanke, 1950) Potonié et Kremp, 1954

Illinites elegans Kosanke, 1950 - Pl. V, fig. 8

Illinites kosankei sp. nov. - Pl. V, fig. 10 (holotype)

Illinites unicus Kosanke, 1950 - Pl. V, fig. 6

Genus *VITREISPORITES* (Leschik, 1955) Jansonius, 1962

Vitreisporites sp. - Pl. V, fig. 12

Genus *JUGASPORITES* (Leschik, 1956) Foster, 1983 emend. Tiwari et Singh, 1984

Jugasporites delasaucei (Potonié et Klaus, 1954) Leschik, 1956 - Pl. V, fig. 4

Jugasporites latus (Leschik, 1956) Foster, 1983 - Pl. V, fig. 7

Jugasporites lueckoides Klaus, 1963 - Pl. V, fig. 9

Jugasporites paradelasaucei Klaus, 1963 - Pl. V, fig. 15

Jugasporites parvus (Klaus, 1963) Foster, 1983 - Pl. V, fig. 5

Jugasporites purus (Leschik, 1956) Tiwari et Singh, 1984 - Pl. V, fig. 11

Jugasporites schaubergeroides Klaus, 1963 - Pl. V, fig. 13

Genus *TRIADISPORA* (Klaus, 1964) sensu Brugman, 1979

Triadispora crassa Klaus, 1964 - Pl. V, fig. 17, 19

Triadispora plicata Klaus 1964 - Pl. V, fig. 16

Triadispora visscheri (Visscher, 1966) comb. nov. - Pl. V, fig. 14

Infraturma *Disaccimonoleti* Klaus, 1963

Genus **LIMITISPORITES** (Leschik, 1956) Klaus, 1963

Limitisporites leschiki Klaus, 1963 - Pl. V, fig. 20

Limitisporites moersensis (Grebe, 1957) Klaus, 1963 - Pl. V, fig. 18

Limitisporites cf. parvus Klaus, 1963 - Pl. VI, fig. 2

Limitisporites rectus Leschik, 1956 - Pl. V, fig. 21

Genus **GARDENASPORITES** Klaus, 1963

Gardenasporites heisseli Klaus, 1963 - Pl. VI, fig. 3

Gardenasporites leonardii Klaus, 1963 - Pl. VI, fig. 1

Gardenasporites cf. moroderi Klaus, 1963 - Pl. V, fig. 22

Gardenasporites cf. oberrauchi Klaus, 1963 - Pl. V, fig. 23

Genus **CHORDASPORITES** Klaus, 1963

Chordasporites sp. - Pl. VI, fig. 13

Subturma **Polysaccites** Cookson, 1947

Infraturma **Polysacciti** Cookson, 1947

Genus **CRUSTAESPORITES** Leschik, 1956

Crustaesporites globosus Leschik, 1956 - Pl. VI, fig. 4

Turma **Plicates** Naumova, 1937

Subturma **Praecolpates** Potonié et Kremp, 1954

Infraturma **Praecolpati** Potonié et Kremp, 1954

Genus **PAKHAPITES** Hart, 1965

Pakhapites sp. - Pl. VI, fig. 5

Subturma **Polysaccites** Erdtman, 1952

Infraturma **Polyplicati** Erdtman, 1952

Genus **GNETACEPOLLENITES** (Thiergart, 1938) Jansonius, 1962

Gnetaceaepollenites steevesi Jansonius, 1962 - Pl. VI, fig. 6

Subturma **Moncolpates** (Wodehouse, 1935) Iversen-Troels et Smith, 1950

Infraturma *Inorti* (Naumova, 1937) Potonié, 1958

Genus *CYCADOPITES* (Wodehouse, 1935) Wilson et Webster, 1946

Cycadopites coxii Visscher, 1966 - Pl. VI, fig. 7

Cycadopites follicularis Wilson et Webster, 1946 - Pl. VI, fig. 8

Genus *MONOSULCITES* (Cookson, 1947) Couper, 1953

Monosulcites sp. - Pl. VI, fig. 9

Group Acritarcha

Genus *LEIOSPHAERIDIA* (Eisenack, 1938) Dovnie, Evitt et Sarjeant, 1963

Leiosphaeridia sp. - Pl. VI, fig. 11

Genus *BALTISPHAERIDIUM* (Eisenack, 1958) Dovnie, Evitt et Sarjeant, 1963

Baltisphaeridium debilispinum Wall et Downie, 1963 - Pl. VI, fig. 15

Baltisphaeridium longispinosum (Eisenack, 1931) Eisenack, 1969 - Pl. VI, fig. 27

Genus *MICHRYSTRIDIUM* (Deflandre, 1937) Sarjeant, 1967

Michrystridium recurvatum Valensi, 1953 - Pl. VI, fig. 12

Michrystridium setasessitante Jansonius, 1962 - Pl. VI, fig. 14

Michrystridium cf. *stellatum* Deflandre, 1945 - Pl. VI, fig. 10, 22, 25

Genus *VERYHACHIUM* (Deunff, 1954) Deunff, 1958 emend. Loebil et Tappan, 1976

Veryhachium irregulare Jekhowsky, 1961 - Pl. VI, fig. 17-19

Veryhachium trispinoides (Jekhowsky, 1961) comb. nov. - Pl. VI, fig. 21

DESCRIPTION OF THE NEW SPECIES, NEW FORMS AND SOME NEW COMBINATIONS

Laevigatisporites giganteus Dybova et Jachowicz, 1957 f. *microsignus* f. nov.

(Pl. I, fig. 5)

Holotype: Pl. I, fig. 5.

Type horizon: Zechstein (PZ1)

Type locality: borehole Siodła IG 1, depth 182,5 m, Holy Cross Mts, Poland.

Derivation of name: [Greek.] *micros* - small, [Lat.] *signum* - mark, a spore with a comparatively small tetrad mark.

Diagnosis. - Amb circular or elliptical with small triradiate tetrad mark. Exine smooth.

Description. - Spore diameter 80-130 μm . Amb circular or elliptical. Arms of distinct tetrad mark straight, 1/4 spore radius long, Exine thin, smooth.

Remarks. - The most characteristic feature is a small size of tetrade mark which differs it from the other forms of this species.

Occurrence. - Poland: the Holy Cross Mountains, Zechstein.

Lycospora permica (Schwartzman, 1976) comb. nov.

(Pl. I, fig. 12)

1976 *Lycospora pseudohirta* var. *permica* sp. et var. nov; E. G. Schwartzman in; J. K. Inosova, A. M. Krusina, E. G. Schwartzman, p. 63, pl. 4, fig. 23, 24.

Holotype: Schwartzman 1976, pl. 4, fig. 23.

Type horizon: Lower Permian.

Type locality: Donieck Basin, Ukraina.

Description. - Spore diameter 32-48 μm . Amb triangular with slightly convex sides. Triradiate tetrad mark distinct with arms narrowed at the ends, reaching outer spore margin. Wide cingulum which becomes transparent toward the equator. Exine proximately fine-grained, on the distal side and cingulum - coarse-grained or pillate.

Occurrence. - Poland: the Holy Cross Mts, Zechstein (PZ1); Ukraina: Donieck Basin, Lower Permian.

Lueckisporites virkkiae Potonié et Klaus, 1954 with a "dark body" f. nov.

(Pl. VII, fig. 7)

1970 *Parmasporites* gen. nov.; H. Kotańska, p. 153-158, pl. 1, fig. 1-3.

Holotype: Pl. VII, fig. 7.

Type horizon: Zechstein (PZ3).

Type locality: borehole Podgace IG 1, depth 97,7 m, the Holy Cross Mts, Poland.

Diagnosis. - Bisaccate pollen grain. Central body outline circular, sacci hemispherical. In the polar position of pollen grain, dark shield ("body") is visible in middle of central body.

Description. - Total grain size 50-80 μm . Bisaccate pollen grain diploxylonoid in outline. Central body circular or ovale. Sacci semicircular or less than semicircular. Sacci bases convex. Dimension of distal zone is of 1/2 central body length. Monolete mark on proximal pole. Exine of central body fine reticulate divided by clearly visible laesura into two hemispheres. In the middle of central body round or oval dark shield of variable size (usually its diameter equals about 1/3 central body diameter), devoided of sculpture. Exine of sacci reticulate.

Remarks. - The specimen is a specific preservation form of *Lueckisporites virkkiae* in the environment with increased salinity. The "dark body" is a preserved, inner cell's contents according to Dybova-Jachowicz. This suggestion can be confirmed by the fact, that the morphological variability, the same as within the *Lueckisporites virkkiae* palynodeme, is observed here. For example *L. virkkiae* norm Aa, Ab, Ac and Bb with a "dark body" were identified.

Occurrence. - Poland, Zechstein, specially numerous in the PZ3 series.

Lunatisporites microsaccatus (Jansonius, 1962) comb. nov.

(Pl. III, fig. 17)

1962 *Taeniaesporites* sp. U; J. Jansonius, p. 65, pl. 14, fig. 1-2.

Holotype: Jansonius 1962, pl. 14, fig. 1.

Type horizon: Lower Triassic.

Type locality: Western Canada.

Derivation of name: [Greek.] *micros* - small, [Lat.] *saccus* - bladder, a pollen grain with the small, reduced bladders.

Description. - Total grain size 15-30 μm . Bisaccate pollen grain haploxylonoid in outline. Central body circular or oval. Sacci folded under central body or strongly reduced so only the distal sacci bases are visible which together with the proximal central taeniae cause an characteristic "check" pattern. Sometimes sacci larger, falciform in outline. Monolete mark at the proximal pole sometimes visible. Exine on proximal side of central body divided

into four taeniae of smooth or fine-grained sculpture. Sacci exine infrareticulate.

Remarks. - The small size of sacci distinguishes this specimen among the other representatives of *Lunatisporites* genus.

Occurrence. - Poland: the Holy Cross Mts, Upper Zechstein- Middle Buntsandstein (PZ3-Tp2); Western Canada, Lower Triassic; Canadian Arctic Archipelago, Lower Triassic.

Illinites kosankei sp. nov.

(Pl. V, fig. 10)

Holotype: Pl. V, fig. 10.

Type horizon: Zechstein (Pzt).

Type locality: borehole Siodła IG 1, depth 183,2 m, Holy Cross Mts, Poland.

Derivation of name: from the name of American palynologist R. M. Kosanke, creator of the genus *Illinites*.

Diagnosis. - Bisaccate pollen grain haploxylonoid in outline. Central body circular or oval. Small tetrade mark. Exine of central body smooth.

Description. - Central body length 32-38 μm (holotype - 36 μm),
central body width 22-30 μm (holotype - 26 μm),
sacci length 20-28 μm (holotype - 26 μm),
sacci width 20-24 μm (holotype - 21 μm).

Bisaccate pollen grain haploxylonoid in outline. Central body oval with longitudinal elongation in polar view and circular in lateral view. Sacci smaller than central body, hemispherical, grouped on the narrow distal side which width equals 1/10-1/8 of central body width. Sometimes sacci touch themselves. Sacci bases convex. Small, triradiate tetrade mark with arms of the 1/6 body radius length on proximal pole. Exine of central body smooth to infragranulate. Sacci exine regular, reticulate.

Remarks. - A small tetrade mark and a very narrow distal side distinguish this species within the other representatives of genus *Illinites*.

Occurrence. - Poland: the Holy Cross Mts, Upper Zechstein.

Triadispora visscheri (Visscher, 1966) comb. nov.

(Pl. V, fig. 14)

1966 *Triadispora muelleri* (Reinhardt et Schmitz, 1962 emend. Reinhardt, 1964) comb.

nov.; H. Visscher, p. 352-353, pl. 12, fig. 1-2, text-fig. 22e.

Holotype: Visscher 1966, pl. 12, fig. 1.

Type horizon: Lower Triassic.

Type locality: Hengelo, the Netherlands.

Derivation of name: from the name of the Dutch palynologist H. Visscher, creator of this species.

Disscusion. - The author proposes to introduce the new species name for the pollen described by Visscher as *Triadispora mulleri* to differentiate it from the monosaccite pollen grain *Hexasaccites muelleri* (Reinhardt et Schmitz, 1962) Reinhardt, 1964.

Description. - Trisaccate pollen grain. Central body big, circular. Sacci less than hemisphaerical in outline, concentrated together on distal side. Triradiate tetrade mark with arms of the 1/4 body radius length on proximal pole. Exina on proximal side of central body granulate or reticulate. Sacci exina delicate, reticulate. Remarks. - The always present three sacci are the most characteristic feature of this species.

Occurrence. - Poland: the Holy Cross Mts, Lower Zechstein; the Netherlands, Lower Triassic.

PALYNOSTRATIGRAPHY

The palynostratigraphic framework established for the Zechstein-Lower Buntsandstein in the Holy Cross Mts is composed of four spore-pollen assemblages which represent the Late Permian Lueckisporites virkkiae and Early Triassic LT1 zones. These assemblages, in ascending stratigraphic order, are:

- I. Lueckisporites virkkiae Ab,
- II. Lueckisporites virkkiae Ac,
- III. Lueckisporites virkkiae Bc,
- IV. Protohaploxylinus sp. div. and Denoisporites playfordii.

These assemblages are described below and summarized in Table 2. Detailed palynozonation of the four selected, most complete boreholes is given in Appendix 1. All taxa defined in this study are alphabetically listed in Appendix 2. The morphology of the selected, age-significant species is described in Appendix 3.

The following criteria were used by the identification of these assemblages:

- 1) the length of the species vertical ranges with a particular allowance for the moment of their first appearance and decline;
- 2) the changes in the frequency of the morphological norms within the *Lueckisporites* palynodeme; the regularity of the lower norms (Aa, Ab, Ba) domination within the older Zechstein deposits, noticed by Visscher (1971, 1978) has been confirmed here;
- 3) the changes in the frequency of monosaccites pollen grains; their quantity decreases in the younger beds;
- 4) the presence of acritarchs.

ASSEMBLAGE I: Lueckisporites virkkiae Ab

The base of this assemblage is defined by the first appearance of the Late Permian taxa with the index species *Lueckisporites virkkiae*, the top - by the last appearance of such species as: *Hamiapollenites cf. bifurcatus*, *Crustaeспорites globosus* and *Plicatipollenites indicus*.

It was identified in the oldest Zechstein deposits, which belong to the PZ1 cycle in the majority of the studied boreholes. It can be divided into three subassemblages: Ia - L. virkkiae Ab and acritarchs, Ib - L. virkkiae Ab and Ic - L. virkkiae Ab and *Strotersporites* sp. div.

SUBASSEMBLAGE Ia: Lueckisporites virkkiae Ab and acritarchs

CHARACTERISTICS - The base of this subassemblage coincides with the assemblage base and the top is defined by the disappearance of acritarchs, which make up 7% of the whole spectrum, and also by the last appearance such species as: *Laevigatisporites minimalis f. pulla*, *Crucisaccites cf. latisulcatus* or *Florinites* sp.

Among the miospores the pollen grains make up 91,6% of the spectrum (in this: Disaccites - 73,6%, Monosaccites - 14,0%, Praecolpates - 3,1% and Polysaccites - 0,8%) and spores - 8,4%.

The disaccate pollen are dominated by *Lueckisporites virkkiae* (19,0% of spectrum) characterized mainly by the low Aa and Ab norms (67,9% of all species). The share of the higher norms accounts respectively: Ac - 3,2%; Ba, Bb - 15,8%; Bc - 4,7%; C - 7,9% and % - 0,5%.

ZECHSTEIN										BUNTSAND		LITHOSTRATIGRAPHY (after Kowalczewski, Rup, 1989; Kuleta, 1990; Zbroja, 1990)		PALYNOMORPHS	
Pz1	Pz2	Pz3	Pzt	-STEIN											
T1	T1r	A1g	T2r	T3	Ca3	A3									
-	-	-	-	-	-	-	-	-	-	<i>Florinites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Pillaspora</i> cf. plurigenus					
-	-	-	-	-	-	-	-	-	-	<i>Vitrisporites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Hamiapollenites</i> cf. bifurcatus					
-	-	-	-	-	-	-	-	-	-	<i>Crustasporites globosus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Triadispora viesscheri</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cordaitina uralensis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Vittatina vittifer</i>					
-	-	-	-	-	-	-	-	-	-	<i>Limitisporites leschikii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Nuskisporites dulhuntyi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Limitisporites moersensis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> NAA, Ab					
-	-	-	-	-	-	-	-	-	-	<i>Klausipollenites schaubergeri</i>					
-	-	-	-	-	-	-	-	-	-	<i>Perisaccus granulatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Illinites unicus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Limitisporites</i> cf. <i>parvus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites delascacei</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites hexagonalis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Vittatina subsaccata</i>					
-	-	-	-	-	-	-	-	-	-	<i>Falcisporites zapfei</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cycadopites coxi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites nouiaulensis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Platysaccus papilionis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Platysaccus niger</i>					
-	-	-	-	-	-	-	-	-	-	<i>Calamospora plicata</i>					
-	-	-	-	-	-	-	-	-	-	<i>Laevigatosporites minimalis</i> f. <i>pulpa</i>					
-	-	-	-	-	-	-	-	-	-	<i>Crucisaccites</i> cf. <i>latisulcatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protohaploxylinus rhomboformis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Klausipollenites decipiens</i>					
-	-	-	-	-	-	-	-	-	-	<i>Gardenasporites</i> cf. <i>moroderi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Pakkhipites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Apiculatisporites apiculatus</i> f. <i>media</i>					
-	-	-	-	-	-	-	-	-	-	<i>Plicatipollenites indicus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites latius</i>					
-	-	-	-	-	-	-	-	-	-	<i>Vesicaspora schemali</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites parvus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Laevigatosporites giganteus</i> f. <i>microsignus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Vittatina costabilis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Gardenasporites</i> cf. <i>oberrauchi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Laevigatosporites vulgaris</i> f. <i>minor</i>					
-	-	-	-	-	-	-	-	-	-	<i>Potoniellisporites simplex</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cordaitina donetziana</i>					
-	-	-	-	-	-	-	-	-	-	<i>Trizonasporites grandis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protohaploxylinus latissimus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Striatopodocarpites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites acutus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites alatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites articolat</i>					
-	-	-	-	-	-	-	-	-	-	<i>Paravesicaspora splendens</i>					
-	-	-	-	-	-	-	-	-	-	<i>Illinites elegans</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites purus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites rectus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Gardenasporites heisseli</i>					
-	-	-	-	-	-	-	-	-	-	<i>Chordasporites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> Bb					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> Bc					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> NC					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> NE					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites paradelasacaei</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protohaploxylinus samoilo维奇i</i>					
-	-	-	-	-	-	-	-	-	-	<i>Endosporites hexarecticulatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Vittatina hiltonensis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Gardenasporites leonardii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Nuskisporites klausii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> NAc					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites lobaccus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lycospora pernica</i>					
-	-	-	-	-	-	-	-	-	-	<i>Verrucosporites pseudomorulae</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites transversundatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites obex</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites gracilis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Klausipollenites stolpini</i>					
-	-	-	-	-	-	-	-	-	-	<i>Klausipollenites minimus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites multiplex</i>					
-	-	-	-	-	-	-	-	-	-	<i>Triadispora crassa</i>					
-	-	-	-	-	-	-	-	-	-	<i>Triadispora plicata</i>					
-	-	-	-	-	-	-	-	-	-	<i>Gnetacepollenites stevensi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lueckisporites virkkiae</i> f. with a "dark body"					
-	-	-	-	-	-	-	-	-	-	<i>Striatobactites ayungii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Striatobactites balmi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protosacculina</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Strepterosporites wilsoni</i>					
-	-	-	-	-	-	-	-	-	-	<i>Strepterosporites richieri</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protohaploxylinus jacobi</i>					
-	-	-	-	-	-	-	-	-	-	<i>Platysaccus leschikii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lunatisporites microsaccatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Illinites kosankei</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites schaubergeroides</i>					
-	-	-	-	-	-	-	-	-	-	<i>Jugasporites lueckoides</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lundbladispora</i> cf. <i>obsoleta</i>					
-	-	-	-	-	-	-	-	-	-	<i>Lundbladispora brevicalyx</i>					
-	-	-	-	-	-	-	-	-	-	<i>Densosporites playfordii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Punctatisporites triassicus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cyclotriletes microgranifer</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cyclotriletes oligogranifer</i>					
-	-	-	-	-	-	-	-	-	-	<i>Guttatisporites</i> sp.					
-	-	-	-	-	-	-	-	-	-	<i>Endosporites papillatus</i>					
-	-	-	-	-	-	-	-	-	-	<i>Protohaploxylinus pantii</i>					
-	-	-	-	-	-	-	-	-	-	<i>Cycadopites follicularis</i>					
-	-	-	-	-	-	-	-	-	-	<i>Monosulcites</i> sp.					
-	-	-	-	-	-	-	-	-	-	MICROPHYTOPLANKTON					
Lueckisporites virkkiae										PALYNOLOGICAL ZONES					
a	b	c	a	b	m					SPORE-POLLEN ASSEMBLAGES AND SUBASSEMBLAGES					

b.c. - basal conglomerate

A 1. 2. 3. 4. 5. 6. 7. 8. 9. 10.

ΣΣ 11. B — a. — b. — c. — d.

Tah 2

Sporomorphs occurrence in the Primo-Triassic sequence in the Holy Cross Mts. A Lithology: 1 - conglomerates, 2 - coarse sandstones, 3 - fine sandstones, 4 - mudstones and siltstones, 5 - limestones, 6 - marly limestones, 7 - oolitic limestones, 8 - dolomites, 9 - marly dolomites, 10 - carbonate nodules, 11 - sulphates. B Palynology: occurrence of sporomorphs: a - sporadically, b - singly, c - not frequent, d - frequent; spore-pollen assemblages: Ia - L. virkkiae Ab and acritarchs, Ib - L. virkkiae Ab and *Strotersporites* sp. div., IIa - L. virkkiae Ac and acritarchs, IIb - L. virkkiae Ac, III - L. virkkiae Bc, IV - *Protohaploxylinus* sp. div and D. playfordii

The genus *Lunatisporites* is the second under the consideration of frequency in this subassemblage (16,0%). It is represented mainly by *L. noviaulensis* (82,0% of species). *L. multiplex*, *L. acutus* and *L. hexagonalis* occur less frequently here.

Klausipollenites schaubergeri, *Limitisporites moersensis* and *Jugasporites delasaucei* are very important elements of this spectrum. The following genera are represented numerously in comparison to the younger assemblages: *Hamiapollenites* (3,2% of spectrum), *Vittatina* (2,7%) and *Vitreisporites* (1,9%).

The monosaccate pollens are dominated by *Nuskoisporites* with species: *N. dulhuntyi* and *N. klausii*. The other taxa (*Cordaitina*, *Perisaccus*, *Potonieisporites*, *Trizonaesporites*, *Plicatipollenites*, *Florinites*, *Crucisaccites*) occur seldom or sporadically.

The Polysaccites are represented only by one species *Crustaeisporites globosus*.

The following species are the most important among spores: *Calamospora pedata*, *C. plicata*, *Laevigatisporites* sp. div. and *Lycospora permica*.

Acritarcha are dominated by *Veryhachium* (75,6% of group). *Baltisphaeridium*, *Leiosphaeridia* and *Michrystrium* occur less frequently.

OCCURRENCE. - This subassemblage was stated in the deposits corresponds to the Copper-bearing Shale (T1), Zechstein Limestone (Ca1) (Fig. 2), locally in the carbonate beds within the "basement conglomerate" and lowermost part of the Lower Anhydrite (A1d) in the 14 boreholes.

COMPARISON AND CORRELATION. - German basin: the Late Permian assemblage, similar to described above but containing more *Lueckisporites virkkiae* specimens (about 40%), was reported from the Copper-Bearing Shale and Zechstein Limestone in Western Germany (Leschik, 1956; Grebe, 1957, Grebe and Schweitzer, 1962) (Tab. 3). Similar subassemblage (EZ1') but with the higher content of acritarchs (24,0%) was described in the Lower Magnesian Limestone, England (Clarke, 1965; Pattison et al., 1973). Alpine basin: *L. virkkiae* Ab and acritarchs subassemblage can be probably correlated with II1 assemblage identified in the upper part of Ad member within the Balaton Red Sandstone Formation, Transdanubian Central Range, Hungary (Barabas-Stuhl, 1990 (unpublished)) and it is similar to the assemblage described from the Grodner Sandstone Formation of Salzgebirges, Austria (Potniw et Klaus; 1954, Klaus, 1955). It can be also correlated with subphase 2 of *L. virkkiae* phase Ab stated in the Val Gardena Formation, Dolomites, Italy (Visscher and Brugman, 1988). The possible correlations with assemblages reported from the

other places outside of Europe are presented in Table 3.

SUBASSEMBLAGE Ib: Lueckisporites virkkiae Ab

CHARACTERISTICS. - The base of the subassemblage is defined by the first appearance of *Lueckisporites virkkiae* f. with a "dark body" and top - by general enrichment of miospores. This is a very poor spectrum in respect both to quality and quantity differentiations of taxa.




Fig.2.

Correlation of the selected, typical Permo-Triassic sequences in the Holy Cross Mts.

Lithology: 1 - conglomerates, 2 - coarse-grained sandstones, 3 - fine-grained sandstones, 4 - sandy mudstones, 5 - mudstones, 6 - siltstones, 7 - limestones, 8 - marly limestones, 9 - marls, 10 - dolomites, 11 - marly dolomites, 12 - carbonate nodules, 13 - sulphates (gypsum)

and anhydrite). Palynology: 14 - spore-pollen assemblages: Ia - *Lueckisporites virkkiae* Ab and acritarchs, Ib - *L. virkkiae* Ab, Ic - *L. virkkiae* Ab and *Strotersporites* sp. div., IIa - *L. virkkiae* Ac and acritarchs, IIb - *L. virkkiae* Ac, III - *L. virkkiae* Bc, IV - *Protohaploxylinus* sp. div. and *Densoisporites playfordii*.

Among the miospores the pollen grains make up 94,0% of the spectrum (in this: Disaccites - 91,9% and Praecolpates - 2,9%) and spores - 4,0%.

The disaccate pollens are dominated by *Lueckisporites virkkiae* (34,4% of spectrum) with Aa and Ab norms and the characteristic form with a "dark body" (4,8% of spectrum). *Lunatisporites noviaulensis* also occurs frequently (22,7%). The content of *Klausipollenites schaubergeri* is 13,5%. The representatives of *Limitisporites*, *Platysaccus*, *Gardenasporites* and *Illinites* are found sporadically.

Spores are dominated by *Calamospora* and *Lycospora*.

OCCURRENCE. - This subassemblage was recognized in the Lower Anhydrite deposits (A1d) only in four boreholes.

COMPARISON AND CORRELATION. - It is difficult to make any correlation of this very poor subassemblage.

SUBASSEMBLAGE Ic: *Lueckisporites virkkiae* Ab and *Strotersporites* sp. div.

CHARACTERISTICS. - The base of this subassemblage is defined by the first appearance of the following striatite pollens: *Strotersporites wilsoni*, *S. richteri*, *Striatoabietites balmei*, *S. aytugii* and *Protosacculina* sp. The top coincides with the whole assemblage top.

Among the miospores the pollen grains make up 94,3% of spectrum (Disaccites - 78,1%, Monosaccites - 13,5%, Praecolpates - 2,5%, Polysaccites - 0,2%) and spores - 5,7%.

Lueckisporites virkkiae, represented by the following norms: Aa, Ab (47,3% of species), Ac (9,3%), Ba, Bb (21,0%), Bc (8,2%), C (4,3%), E (0,6%) and form with a "dark body" (9,3%), dominates within spectrum (28,1%). An increase of higher norms frequency is observed here in comparison with the older subassemblages. The specimens of *Klausipollenites* and *Lunatisporites* occur in the same quantity (about 15% of spectrum). This

last genus is represented not only by *L. noviaulensis* but also by: *L. alatus*, *L. labdacus*, *L. ortisei* and *L. gracilis*. The other taxa occur singly or scattered. An increase number of *Falcisporites* specimen (3,4% of spectrum) is remarkable. Distinctly has decreased the frequency of *Vittatina* (to 0,9%), *Vitreisporites* (0,6%) and *Hamiapollenites* (0,3%).

Nuskoisporites is still dominated among the monosaccate pollens. The representatives of *Perisaccus*, *Cordaitina* and *Potonieisporites* are less abundant.

Spores are dominated by *Calamospora* and *Laevigatosporites*.

OCCURRENCE. - This subassamblage was found in the Terrigenous Series (T1r) and Upper Anhydrite (A1g) deposits in majority of studied boreholes. Only in the one borehole its base lies in the uppermost part of the Lower Anhydrite (A1d).

COMPARISON AND CORRELATION. - German basin: Late Permian assemblage, similar to described above, but containing twice more *Klausipollenites* specimen (35% of spectrum) was reported from the Upper Anhydrite of Western Germany (Leschik, 1956). A very big likeness exists between the EZ" subassemblage from the upper part of the Lower Magnesian Limestone, England (Clarke, 1965; Pattison et al. 1973). This last one differs only in the higher (10,9%) content of *Falcisporites*. Alpine basin: This subassemblage can be correlated with the subphase 4 of Ab Lueckisporites phase described in the top of Val Gardena Formation, Italy (Visscher and Brugman, 1988) (Tab. 3).

ASSEMBLAGE II: Lueckisporites virkkiae Ac


The base of this assemblage is defined by the first appearance of *Lunatisporites microsaccatus*. The top boundary is usually difficult for clear determination.

It was recognized in the PZ3 cycle sediments in the eight boreholes. It can be divided into two subassemblages: IIa - *L. virkkiae* Ac and acritarchs and IIb - *L. virkkiae* Ac.

SUBASSEMBLAGE IIa: Lueckisporites virkkiae Ac and acritarchs

CHARACTERISTICS. -The base of this subassemblage coincides with the assemblage base and top is defined by *Jugasporites parvus* and *Vesicaspora schemeli* last appearance and by the disappearance of acritarchs which make up 2,3% of spectrum.

Miospores are dominated by pollen grains (92,8% in this: Disaccites - 71,7%,



Tab. 3.

Correlation of the Late Permian and Early Triassic spore-pollen assemblages.

a - assemblages identified in the Holy Cross Mts; Authors: 1 - Wagner 1987, Wagner et al. 1978, 1981; 2 - Dybova-Jachowicz 1974, 1984a, 1984b, Dybova-Jachowicz et al. 1984, 1987, Jachowicz, Dybova-Jachowicz 1987, Orłowska-Zwolińska 1985; 3 - Kowalczewski, Rup 1989, Rup 1985, Zbroja 1990; 4 - Kuleta 1985, 1990; 5 - Barabas 1990 (unpublished); 6 - Barabas-Stuhl 1990 (unpublished); 7 - Klaus 1955; 8 - Potonié, Klaus 1954; 9 - Massari et al. 1988; 10 - Visscher, Brugman 1988, 11 - Brugman 1983; 12 - Klaus 1963; 13 - Grebe 1957; 14 - Grebe, Schweitzer 1962; 15 - Leschik 1956; 16 - Geiger, Hopping 1968; 17 - Pattison et al. 1973; 18 - Clarke 1965; 19 - Smith, Brunstrom 1974; 20 - Smith, Crosby 1979; 21 - Visscher 1971; 22 - Balme 1979; 23 - Jansonius 1962; 24 - Mc Gregor 1965; 25 - Utting 1987; 26 - Luber, Waltz 1941; 27 - Kiuntzel 1965; 28 - Molin, Koloda 1972; 29 - Gomanikov, Meyen 1986; 30 - Severcova 1966; 31 - Molin 1989; 32 - Grausmann et al. 1989; 33 - Chalyshev et al. 1965; 34 - Chalyshev, Wariukhina 1966, 1968; 35 - Waruikhina 1971; 36 - Inosova et al. 1976; 37 - Sadownikow 1989; 38 - Romanowskaya 1963; 39 - Romanowskaya et al. 1973; 40 - Sadownikow 1989; 41 - Dibner 1958; 42 - Shu 1982, 1986; 43 - Shu, Norris 1988; 44 - Zhi, Jiazheng 1987; 45 - Balme 1970; 46 - Visscher 1967; 47 - Maheshwari, Banieryj 1975; 48 - Srivastava et al. 1988; 49 - Eshet 1990; 50 - Jekhowsky, Goubin 1964; 51 - Goubin 1965; 52 - Foster 1979; 53 - Utting 1979; 54 - Falcon 1973; 55 - de Jersey 1970, 1979; 56 - Kremp et al. 1977; 57 - Foster 1982; 58 - Balme, Hennelly 1955; 59 - Grebe 1970; 60 - Helby 1973; 61 - Balme 1963.

Monosaccites 13,1%, Praecolpates - 5,2% and Polysaccites - 0,5%) and spores make up 7,2% of spectrum.

Lueckisporites virkkiae, represented by the following norms: Ab (20,0% of species), Ac (20,9%), Ba, Bb (28,4%), Bc (12,0%), C (6,3%), E (3,4%) and form with a "dark body" (9,0%), is the most frequent (20,8% of spectrum). *Klausipollenites* with species: *K. schaubergeri*, *K. staplini* and *K. minimus*, occurs in the same number (20,0%). *Lunatisporites* is less frequent (14,1%). The striatiti pollens are comparatively abundant.

The monosaccate pollens are dominated by *Nuskoisporites*. The representatives of *Perisaccus* and *Cordaitina* occur singly.

The acritarchas are represented by *Baltisphaeridium* (40,7% of microplankton), *Veryhachium* (38,1%) and *Michrystridium* (21,2%).

OCCURENCE. - This subassemblage was stated in the deposites correspoding to the Grey Pelite (T3) and Platy Dolomite (Ca3) in eight boreholes.

COMPARISON AND CORRELATION is given below, commonly for IIa and IIb subassemblages.

SUBASSEMBLAGE IIb: Lueckisporites virkkiae Ac

CHARACTERISTICS. -The boundaries of this poor subassemblage are diffucult for determination.

The spectrum is dominated by *Lueckisporites virkkiae* (29,3%). The representatives of *Lunatisporites* make up 18,7% of spectrum and *Klausipollenites* - 12,0%. The other miospores occur singly.

OCCURENCE. - This subassemblage was distinguished in the deposits corresponding to the Main Anhydrite (A3) and in the lowermost part of the Top Terrigenous Series (Pzt) in the three boreholes.

COMPARISON AND CORRELATION. - Alpine basin: Lueckisporites virkkiae Ac assemblage shows a big similarity to subphaze 5 of Lueckirrkkiae Ac phaze described from the Lower Bellerophon Formation, Dolomites, Italy (Visscher, Brugman, 1988).

ASSEMBLAGE III: Lueckisporites virkkiae Bc

CHARACTERISTICS. - The base of this assemblage is determinated by the first appearance of *Jugasporites lueckoides*, *J. schaubergeroides* and *Illinites kosankei*. The top is assigned by declination of the most of the Late Permian species with the index taxa *Lueckisporites virkkiae*.

Pollen grains dominated among miospores (93,9% of spectrum), misopres make 6,1%.

Lueckisporites virkkiae represented by following norms: Ab (22,5% of species), Ac (19,8%), Ba, Bb (22,1%), Bc (25,3%), C (6,6%) and E (1,2%) occur the most frequent within pollens 24,3% of spectrum. Repräsentatives of *Lunatisporites* make up 15,9% of

assemblage. Such species as *L. labdacus*, *L. multiplex* and *L. gracilis* are more common than in the older assemblages. The share of *Klausipollenites* equals 14,5% and *Falcisporites* - 5,4%. The other pollens occur rare or scattered.

Monosaccate pollens are dominated by *Perisaccus*.

The genus *Calamospora* is the most frequent among the spores.

OCCURENCE. - This assemblage was identified in the higher part of the Top Terrigenous Series (Pzt) in the five boreholes.

COMPARISON AND CORRELATION. - Alpine basin: The assemblage which can be correlated to *Lueckisporites virkkiae* Bc was described as assemblage II3 from the top of Balaton Red Sandstone and lower part of evaporate Tabajd Formations, Transdanubian Central Range, Hungary and to assemblage II3 from the Kövágószólös Sandstone formation, Mecsek Mountains, Hungary (Barabas-Stuhl, 1981, 1990 (unpublished)). This last differs in the higher contents of *Jugasporites* (24% of assemblage). There exist also a similarity to the phase *Lueckisporites virkkiae* Bc distinguished by Visscher and Brugman (1988).

ASSEMBLAGE IV: *Protohaploxylinus* sp. div. and *Densoisporites playfordii*

CHARACTERISTICS. - The base of this assemblage is defined by the first appearance of Trassic species: *Lundbladispora brevicula*, *L. cf. obsoleta*, *Densoisporites playfordii*, *Cyclotriletes ologranifer*, *C. microgranifer*, *Punctatisporites triassicus*. The top is assigned by the first appearance of *Densoisporites neburgii*.

Miospores make up 93,7% of spectrum, acritarchs - 5,0% and so called "Tympanicysta" forms - 1,3%.

The pollen grains still dominate among the miospores (80,3%) whereas the rest of spores equals 19,7%.

Dissaccites are the most frequent within the pollens (39,7% of spectrum). They are dominated by striatini specimens from such genera as: *Protohaploxylinus* (18,5% of spectrum) represented mainly by *P. pantii* and *P. samoilovichii*, *Lunatisporites* (13,6%) with *L. noviaulensis*, *L. transversundatus*, *L. microsaccatus*, *Striatoabietites* and *Strotersporites* (8,4%). The share of *Klausipollenites* represented by *K. staplinii* and *K. minimus*, equals 8,5%.

Monosaccate pollens (6,0% of spectrum) are dominated by *Endosporites papillatus*.

Cycadopites coxii, *C. follicularis* and *Gnetacaesporites steevesi* are the most common within Praecolpates.

Spores are dominated by *Densoisporites* (10,0% of spectrum) represented by *D. playfordii*. *Lundbladispora* and *Cyclotriletes* occur less frequent. The other spores are single or scattered.

Acritarchs are represented mainly by: *Baltisphaeridium* (83,2% of acritarchs), *Veryhachium* (12,3%) and *Michrystridium* (4,5%).

OCCURENCE. - This spectrum was recognized in the A1 complex deposits of the Lowermost Buntsandstein in the six boreholes.

COMPARISON AND CORRELATION. - The Lowemost Triassic assemblage is known, under the different names, from several localites in Europe and the world (Tab. 3). German basin: The similar assemblage, as a *Lundbladispora obsoleta-Protohaploxylinus pantii* zone, was described from the Lower Buntsandstain of central Poland (Dachów M-24, Otyń IG 1 and Florentyna 2 boreholes) by Orłowska-Zwolińska (1984, 1985). It differs from the Holy Cross assemblage only by lack of acritarchs. The share of individual miospore instead is almost identical. Alpine basin: The *Protohaploxylinus* sp. div. and *Densoisporites playfordii* assemblage can probably be correlated to II4 assemblage distinguished in the Tabajd Evaporites formation of Transdanubian Central Range and Upper part of Kövágószólós Sandstone formation of Mecsek Mts in Hungary (Barabas-Stuhl, 1981, 1990 (unpublished)). This last differs from the Holy Cross one by the mixed Late Permian-Early Triassic charcter. The Late Permian elements make up 75% of spectrum and are dominated by *Lueckisporites virkkiae* Bc and C. There is also a big similarity to *Striatiti-Triletes-Tympanicysta* assemblage described from the Tessero member of Werfen formation in Dolomites (Massari et al. 1988). *Protohaploxylinus* sp. div. and *Densoisporites playfordii* assemblage belongs to LT1 zone of European, palynological scheme proposed by Brugman (1983).

RELATIONS BETWEEN MICROFLORISTICAL ASSEMBLAGES AND LITHOSTRATIGRAPHY

If the Permo-Triassic assemblges are considered in the chronostratigraphic aspect there arises a problem how far palynologic assotiations are environmentally dependent. In the case

then lithological units reflect their depositional environment, the possibility of treatment the assemblage boundaries as time-line should be proved by their independence from lithostratigraphic boundaries. Table 2 shows that most of assemblage does not coincide with lithological complexes boundaries: the base of *Lueckisporites virkkiae* Ab and acritarchs subassemblage is found in most of the boreholes at the lower part of the Zechstein Limestone (Ca1) but it also includes the underlaying basement conglomerate and locally, Copper-Bearing Shale (T1). The top of this subassemblage is stated in the upper part of Zechstein Limestone and in the lowermost part of the Lower Anhydrite (A1d) deposits. Both boundaries of *Lueckisporites virkkiae* Ab subassemblage lay within the Lower Anhydrites series. The basis of *Lueckisporites virkkiae* Ab and *Strotersporites* sp. div. subassemblage is usually found at the lower part of the recessive Terrigenous Series (T1r) but, in the case of one borehole, also at the uppermost part of the Lower Anhydrite. The top of this subassemblage lies in the upper part of the Upper Anhydrite (A1g) deposits. *Lueckisporites virkkiae* Ac and acritarchs subassemblage is the only which boundaries coincide with the lithological boundaries of the Grey Pelite (T3) and Plate Dolomite (Ca3) complexes. The base of *Lueckisporites virkkiae* Ac subassemblage is stated within the Main Anhydrite (A3) series and top - in the upper part of the Main Anhydrite and lower part of the Top Terrigenous Series (Pzt). And Early Triassic *Protohaploxylinus* sp. div. and *Densoisporites playfordii* boundaries are found within the A1 lithological complex.

The distinguished assemblages have a relatively consistent occurrence through the studied area both in the shallow zone near the Paleozoic core and deeper one placed far off it. These facts entitle to treatment the assemblage boundaries as the time-lines.

There exists also a possibility of drawing a correlation between the Holy Cross assemblages and the Late Permian-Early Triassic spectra known from the other regions of Poland which had represent different zones of the sedimentary basen during the Permian-Triassic period.

The microfloristic assemblages very similar to *Lueckisporites virkkiae* Ab assemblage were recognized in the Lower Zechstein deposits of the Nida basen (south from the Holy Cross Mts) (Dybova-Jachowicz, Laszko, 1978; Fijałkowska, 1990 (unpublished), in the Lower Zechstein in Mielnik borehole (E Poland) (Orłowska-Zwolińska, 1962) and in the Lower Zechstein deposits from afew boreholes on the precambrian platform (NE Poland) (Dybova- Jachowicz, 1981). All these localities together with the Holy Cross Mts area

belonged to the shallow, costal part of the basen. The assemblages which can be correlated to *Lueckisporites virkkiae* Ab spectrum were recognized in the Copper- Bearing Shale and Zechstein Limestone, representing the deeper zones of the basen from the fore-Sudetic monocline (SW Poland) (Kotańska, Krasoń, 1966; Rospondek et al., 1993) and Western Pomerania (NW Poland) (Dybova-Jachowicz, 1981; Jachowicz, 1991).

The assemblages similar to *Lueckisporites virkkiae* Ac were described from the Plate Dolomite on the fore-Sudetic monocline (Kotańska, Krasoń, 1966; Kotańska, 1970; Górecka, Parka, 1980) and in Western Pomerania (Dybova-Jachowicz, 1981).

The spectra which can be correlated to *Lueckisporites virkkiae* Bc were identified within the PZ3-PZ4 deposits from the Upper Silesia basin margin (Dybova-Jachowicz et al., 1991) and from central Poland (Dybova-Jachowicz et al. (in press)).

The correlation with the Early Triassic assemblge has been presented in the former chapter.

BOUNDARY PROBLEM

The precise definition of the Permian-Triassic boundary is still an open question in the Holy Cross Mts in spite of fact that is has been a subject of detailed study for last thirty years. The main reason of this situation is a total luck of fossils remains in the boundary beds. Therefore the determination of this boudnary has been grounded only on the lithological premises so far. In the older works (Senkowiczowa, Ślączka, 1962; Rup, 1985; Kuleta, 1985) the boundary between the Zechstein and Buntsandstein deposits was put at the top of the A0 "knobby" complex inculuded then into the Top Terrigenous Series (Pzt). The Buntsandstein deposits begun from A1 sandy complex. The results of current, sedimentological studies (Pieńkowski, 1987, 1989; Kuleta, 1990) improved the principle of leveling this boundary and including the A0 complex to the Triassic. In that case the Zechstein-Buntsandstein boundary would lay at the top of the Top Terrigenous Series. Unfortunately any sporomorphs has been found so far in the reddish and mottled mainly mudstone deposits of A0 complex and the first Triassic assemblage appeares only in the next A1 unit. There is then a gap in the palynological scheme comprised the upper part of the Top Terrigenous Series, A0 complex and lower part of A1 complex.

PALEOCLIMATE AND PALEOENVIRONMENT

The suitability of sporomorphs for paleoenvironmental interpretations has been unquestionable since Robinson's (1973), Kremp's (1977), Meyen's (1987) and Visscher and Van Der Zwan's (1981) works.

To evaluate Permian-Triassic paleoenvironments in the Holy Cross Mts, the model of Visscher and Van Der Zwan (1981) was applied to the recovered assemblages and subassemblages. The 16 morphological groups of sporomorphs were used in the statistical analysis of assemblage composition. Results are given in the Figure 3, which shows the generalized histograms for all distinguished assemblages and subassemblages.

The significant dominance of the xerophytic elements (Groups H-L) with the striatite pollen grains (Group I) is observed in all assemblages. It indicates generally arid climatic conditions on the studied area during the Late Permian. This suggestion is also confirmed by the lithological data (Wagner et al., 1987; Kowalczewski, Rup, 1989).

Microphytoplankton in the first subassemblage evidences the Zechstein transgression and shallow water conditions during the sedimentation of the Zechstein Limestone. Then the climate became more dry in the Lower Anhydrite period and *Lueckisporites virkkiae* Ab subassemblage contains almost the same xerophytic forms (88,9% of spectrum). There was marked a short-time increase of humidity during the sedimentation of the recessive Terrigenous Series and the dry conditions returned in the Upper Anhydrite which is reflected in the increase of striatite pollens race.

Acritarchs in *Lueckisporites virkkiae* Ac subassemblage evidence the second Zechstein transgression which reached the Holy Cross Mts. A slight increase of humidity is observed in comparison to the older spectra (11,4% of the hygrophytic elements). The approximate conditions prevailed till the end of Permian.

The significant development of the humid vegetation had place only in Early Triassic that can be inferred from the higher race of hygrophytes in *Protohaploxylinus* sp. div. and *Densoisporites playfordii* assemblage (40,9%). The presence of the single acritarchs is connected with the Buntsandstein transgression. This increase of climate humidity in the Early Triassic is mentioned also in the paleoclimatic interpretations from other regions of Europe and of the world (Robinson, 1973; Parris et al., 1982).

The reconstruction of the depositional environment, based on a combination of the modified model for the theoretical distribution of organic remains connected with the sedimentation and sedimentary data (Kuleta, 1985; Rup, 1985), is presented on Figure 4. The inferred depositional conditions seem to be comparatively stable, generally in sebkha environment with the three periods of transgression and deposition in the shallow lagoon (the Copper-Bearing Shale, Zechstein Limestone, Grey Pelite, Platy Dolomite and Early Triassic A1 complex). The Uppermost Permian deposits originated in the playa environment and the Lowermost Triassic sandstones have a character of the river deposits (Kuleta, 1985).




Fig. 3.

Application of the environmental model to the Upper Permian and Lower Triassic samples in the Holy Cross Mts.

1 - hygrofile elements, 2 - mixed elements, 3 - xerofile elements, 4 - microphytoplankton.

A - monolete, acavate spores, B - trilete, acavate, laevigate and apiculate spores, C - trilete, acavate and verrucate spores, D - trilete, cingulate and zonotrilete spores, E - monosulcate pollen, F - alete (proto) bisaccate pollen, G - trilete (proto) bisaccate pollen, H - monolete

(proto) bisaccate pollen, I - taeniae (proto) bisaccate pollen, J - Triadispora-complex, K - vesicate pollen, L - (proto) monosaccate pollen, M - Leiosphaeridia, N - Michrystridium, O - Baltisphaeridium, P - Veryhachium. Spore-pollen assemblages: A - Lueckisporites virkkiae Ab and acritarchs, B - L. virkkiae Ab, C - L. virkkiae AB and Strotersporites sp. div., D - L. virkkiae Ac and acritarchs, E - L. virkkiae Ac, F - L. virkkiae Bc, G - Protohaploxylinus sp. div. and Densoisporites playfordii.




Fig. 4.

Reconstruction of the depositional environment during the Late Permian and Early Triassic in the Holy Cross Mts.

- 1 - big spores,
- 2 - middle-size spores,
- 3 - small spores,
- 4 - big pollen,
- 5 - small pollen,
- 6 - microphytoplankton.

CONCLUSIONS

1. Microflora, as so far, has been a base of the Permian-Triassic biostratigraphy in the Holy Cross Mountains. It allowed to create a comparatively complete palynological scheme which contains three Late Permian and one Early Triassic assemblage. The Late Permian assemblages represent *Lueckisporites virkkiae* zone and Early Triassic one - LT1 zone. These assemblages (from older to younger) are mentioned below.

1) *Lueckisporites virkkiae* Ab assemblage was recovered within the oldest Zechstein deposits (PZ1). It is dominated by *Lueckisporites virkkiae* norm Ab specimens. Such genera as *Hamiapollenites*, *Vittatina* and *Crustaeспорites* are characteristic here. It can be divided into three subassemblages: *L. virkkiae* Ab and acritarchs which occurs in the "basement conglomerate", Copper-Bearing Shale, Zechstein Limestone and lowermost part of the Lower Anhydrite; *L. virkkiae* Ba - in the upper part of the Lower Anhydrite and *L. virkkiae* Ab and *Strotersporites* sp. div. - in the Terrigenous Series and Upper Anhydrite. This last subassemblage is distinguished by the higher contents of the striatite pollens.

2) *Lueckisporites virkkiae* Ac assemblage was identified within the PZ3 deposits. It is dominated by Ac norm of *Lueckisporites virkkiae*. The striatite pollens belonged to *Lunatisporites* and *Strotersporites* genera are abundant here. This assemblage can be divided into two subassemblages: *L. virkkiae* Ac and acritarchs which occurs in the Grey Pelite and Platy dolomite and *L. virkkiae* Ac - in the upper part of the Main Anhydrite and Lowermost part of the Top Terrigenous Series.

3) *Lueckisporites virkkiae* Bc assemblage was recognized within the youngest Permian deposits in the upper part of the Top Terrigenous Series. It is characterized by the high frequency of *Lueckisporites virkkiae* Bc norm and striatite pollens.

4) *Protohaploxylinus* sp. div. and *Densoisporites playfordii* assemblage occurs in the Lower Triassic A1 complex. It is characterised by the first appearance of Triassic taxa as *Densoisporites*, *Lundbladispora* or *Cyclotriletes*.

The distinguished assemblages have a relatively constant position in the section on the studied area. Most of their boundaries does not coincide with the lithological boundaries. These facts let treat them as a time-lines.

Unfortunately any sporomorphs were found at the Zechstein\Buntsandstein boundary thus the distinguishing between them still has to be based on the sedimentary criteria.

One should remember, that presented schema has a local character due to geological conditions. The Zechstein-Buntsandstein section contains namely numerous stratigraphical gaps in the Holy Cross Mts.

2. Spore-pollen assemblages, identified in the Holy Cross Mts can be correlated to the contemporaneous spectra referred from the other regions of Europe assigned both to the German basin (Germany, North Sea Basin, England, Ireland) and to Tetyd (Hungary, Austria, Italy). There is also a similarity to the assemblages known from the other continents.

3. Microflora precisely reflects the paleoclimatic changes during the individual Zechstein cyclothemes. It confirms the suggestion about the arid climate obtained from the sedimentary and lithological studies. An increase of humidity is observed only in the Early Triassic.

4. Microflora is also good indicator of the depositional environment. It can be inferred on its base that the sedimentation in sebkha environment dominated during the Late Permian with few periods of the deposition in the deeper conditions of shallow lagoon.

ACKNOWLEDGMENT

This study is based on the author's Ph. D. thesis conducted at the State Geological Institute of Kielce and supervised by Prof. dr. hab. S. Dybova-Jachowicz (State Geological Institute of Sosnowiec). It was reviewed by Dr. hab. E. Turnau (Polish Academy of Sciences, Cracow) and Prof. dr. hab. Z. Kotański (State Geological Institute of Warsaw).

Special thank to promotor Prof. dr. hab. S. Dybova-Jachowicz, Direction of State Geological Institute at Warsaw and members of State Geological Institute at Kielce: Dr. hab. Z. Kowalczewski, St. Zbroja, M. Kuleta and M. Rup.

REFERENCES

- BALME B.E., 1963 - Plant microfossils from the Lower Triassic of Western Australia. *Palaeontology*, 6/1, pp. 1-40.
- BALME B.E., 1970 - Palynology of Permian and Triassic Strata in the Salt Range and Surghar Range, West Pakistan. In: B. Kummel, C. Teichert (Eds.): *Stratigraphic boundary problems; Permian and Triassic of West Pakistan*. Univ. Kansas Spec. Publ., 4, pp. 360-453.
- BALME B.E., 1979 - Palynology of Permian-Triassic boundary beds at Kap Stosch, East Greenland. *Medd. Gronland*, 200/6, pp. 1-37.
- BALME B.E., HENNELLY J.P.F., 1955 - Bisaccate sporomorphs from Australian Permian Coals. *Aust. J. Bot.*, 3/1, pp. 89-98.
- BALME B.E., HELBY R.J., 1973 - Floral modification at the Permian- Triassic boundary in Australia. In: A. Lohgan, L. V. Hills (Eds.): *The Permian-Triassic systems and their mutual boundary*. Can. Soc. Petrol. Geol. Mem., 2, pp. 433-444.
- BARABAS-STUHL F., 1981 - Microflora of the Permian and Lower Triassic sediments of the Mecsek Mountains (S. Hungary). *Acta Geol. Acad. Sci. Hung.*, 24/1, pp. 49-97.
- BRUGAMAN W.A., 1983 - Permian-Triassic Palynology. State Univ. of Utrecht, 121 p.
- CHALYSHEV W.I., WARIUKHINA L.M., MOLIN W.A., 1965 - Permian - Triassic boundary in the red bed deposits of the Norhtern Preural. Nauka, Moscow, pp. 115 (in Russian).
- CHALYSHEV W.I., WARIUKHINA L.M., 1966 - Biostratigraphy of the Trassic deposits in Pechora Basin. Kom. Fil. Inst. Geol. Akad. Nauk SSSR, Moscow, pp. 145 (in Russian).
- CHALYSHEV W.I., WARICHINA L.M., 1968 - The Upper Permian stratigraphy of the north-eastern European part of the USSR. Nauka, Leningrad, pp. 244 (in Russian).
- CLARKE R.F.A., 1965 - British Permian Saccate and Monosulcates miospores. *PALAEONTOLOGY*, 8/2, pp. 322-354.
- CZARNIECKI S., KOSTECKA A., KWIATKOWSKI S. (1965) - *Horridonia horrida* (Sowerby) from the Zechstein Conglomerate of Gałezice area. *Rocznik Pol. Tow. Geol.*, 35/4, pp. 467-473, Cracow.

- DIBNER A.F., 1958 - On the age of the Upper Paleosoic deposits of the Nordvik region according to palynological data. *Trudy NIIG Arkt.*, 67/79, pp. 73-85, Moscow (in Russian).
- DIBNER A.F., 1971 - Pollen of Cordaites from Angarida. *Trudy VNIGRI, Palaeont. and Biostrat. Ser.*, 32, pp. 77, Leningrad (in Russian).
- DYBOVA-JACHOWICZ S., 1974 - Analyse palinologique des sediments rouges salifères du Zechstein Supérieur ("Zouber Rouge") à Kłodawa, Pologne. *Rev. Palaeobot. and Palynol.*, 17, pp. 57- 61.
- DYBOVA-JACHOWICZ S., 1976 - From the study upon the Zechstein microflora in the Holy Cross region. In: *Mat. 2nd Paleontol. Confer.*, Kielce, 14-16th of October 1976, pp. 11-12, (in Polish).
- DYBOVA-JACHOWICZ S., 1981 - The palynological assemblage from Upper Permian of Poland. *Proc. Inter. Symp. Central. Europ. Permian*, Jabłonna 27-29th April, 1978, Warsaw, pp. 456-470.
- DYBOVA-JACHOWICZ S., 1984a - Palynology of the Permian-Triassic boundary in Poland. *Abstr. 6th Inter. Palynol. Conf.*, Calgary 1984, pp 37.
- DYBOVA-JACHOWICZ S., 1984b - Results of a palynological investigation of the Uppermost Carboniferous and Permian of Poland. *Ibidem*, pp. 38.
- DYBOVA S., JACHOWICZ A., 1957 - Microspores of the Uppersilesian Productive Carboniferous. *Pr. IG*, 23 (in Polish).
- DYBOVA-JACHOWICZ S., JAGIELSKA L., 1977 - Acritarchs occurrence in the Holy Cross Zechstein. *Pr. Nauk. Uniwersyteckie Śl.*, 192, pp. 99- 113 (in Polish).
- DYBOVA-JACHOWICZ S., LASZKO D., 1976 - The characteristics of spore-pollen spectrum form the Permian-Triassic boundary beds in the Holy Cross Mts. In: *Mat. 2nd Paleontol. Confer.*, Kielce, 14-16th of October 1976, pp. 23-24 (in Polish).
- DYBOVA-JACHOWICZ S., LASZKO D., 1978 - Spore-pollen spectrum of the Permian-Triassic junction beds in Holy Cross Mts. In: R. Wagner, G. Piątkowski (Eds.): *Guide of Ecxsurs. part 2, Symp. on Europaen Central Permian*, pp. 81-106, Warsaw.
- DYBOVA-JACHOWICZ S., LASZKO D., 1980 - Spore-pollen spectrum of the Permian and Triassic deposits from the Piekoszów syncline in the Holy Cross Mountains. *Kwart. Geol.*, 24/3, pp. 611-641 (in Polish).

- DYBOVA-JACHOWICZ S., ORŁOWSKA-ZWOLIŃSKA T., WAGNER R., 1984 - Permian-Triassic boundary in the Polish depositional basin (litho- and palynostratigraphical results). Abstr. 27th Inter. Geol. Congr., Moscow 1984. pp. 87-92.
- DYBOVA-JACHOWICZ S., POKORSKI J., 1984 - The Stratigraphy of the Carboniferous and Lower Permian in Strzeżewo IG 1 borehole. Kwart. Geol., 28/3-4, pp. 589-616 (in Polish).
- DYBOVA-JACHOWICZ S., POKORSKI J., WAGNER R., 1987 - Litho- and palynological results of investigations of Carboniferous-Permian and Permian-Triassic boundaries in Poland and their correlation with the selected profiles. In: Abstr. 11th Inter. Carbon. Congr. in China.
- DYBOVA-JACHOWICZ S., KIERSNOWSKI H., TRZEPIERCZYŃSKA A., 1991 - Upper Carboniferous of the Upper Silesian Margin. In: Mat. Symp. Geology of the coal-bearing formations, Cracow, pp. 13-18 (in Polish).
- ESHET Y., 1990 - Paleozoic-Mesozoic palynology of Israel. 1. Palynological Aspects of the Permo-Triassic Succession in the Subsurface of Israel. Geol. Surv. of Israel Bull. 81, pp. 1-57.
- FALCON R.M.S., 1973 - Palynology of the Lower Karroo Succession in the middle Zambezi basin. Bull. Geol. Surv. N. Dhod., 70, pp. 43-71.
- FOSTER C.B., 1979 - Permian plant microfossils of the Blair Athol Coal Measures, Baralaba Coal Measures and Basen Revan Formation of Queensland. Geol. Surv. Queens. Publ., 374, Palaeont. Pap. 45, 244 p.
- FOSTER C.B., 1982 - Spore-pollen assemblages of the Bowen Basin, Queensland (Australia): Their relationship to the Permian/ Triassic boundary. Rev Palaebot. and Palynol. 36, pp. 165- 183.
- FOSTER C.B., 1983 - Jugaporites Leschik 1956 a Late Palaeozoic operculate pollen genus. Mem. Ass. Austral. Palaeontol. 1, pp. 327-338.
- GEIGER M.E., HOPPING M.C., 1968 - Triassic stratigraphy in the Southern North Sea Basin. Phil. Trans. Roy. Soc. London, Ser. 750/254, pp. 1-36.
- GOMANIKOW A.W., 1983 - Palaeobotanical characteristic of the Upper Tatarian of the Russian Platform. Akad. Nauk. SSSR., Geol. Inst., Moscow (in Russian).

- GOMANIKOW A.W., MEYEN S.V., 1986 - Tatarian flora. Trudy Izdat. Nauka, 401, Moscow, 174 p .
- GOUBIN N., 1965 - Description et repartition des principaux pollenites permies, triassiques et jurassiques des sondages du Basin de Morondava (Madagaskar). Rev. Inst. Franc. Petrole Ann. Combust. Liquides 20/10, pp. 1415-1461.
- GÓRECKA T., PARKA Z., 1980 - Results of the palynological investigations of the Permo-Carboniferous towards NE from Wrocław. Pr. Nauk. Inst. Górn. 35/16, Wrocław, pp. 1-33 (in Polish).
- GRAUSMANN W.W., TUMANOWA I.J., 1989 - Palaeontological characteristics of the Lower Triassic deposits in the north- western part of the Wiliusk syncline. Tr. Inst. Geol. and Geogr. Acad. Nauk SSSR 722, Moscow, pp. 50-52 (in Russian).
- GREBE H., 1957 - Zur Mikroflora des niederrheinischen Zechstein. Geol. Jb. 73, pp. 51-74.
- GREBE H., 1970 - Permian plant microfossils from the Newcastle Coal Measures (Namabeen Group boundary, Lake Munmorah, New South Wales). Rec. Geol. Surv. New South Wales 12, pp. 125- 136.
- GREBE H., SCHWIETZER H.J., 1962 - Die Sporae dispersae des niederrheinischen Zechstein. Fortschr. Geol. Rheinland Westf. 12, pp. 201-224.
- HART G.F., 1964 - A review of the classification and distribution of the Permian miospore: Disaccate Striatiti. Abstr. 5th Inter. Congr. Stratigraphy and Geology of Carboniferous, Paris 9-12th Sept. 1963, pp. 1171-1199.
- HART G.F., 1965 - The systematic and distribution of Permian miospores. Witwatersrand Univ. Press, Johanesburg, 252 p.
- HELBY R.J., 1973 - Review of Late Permian and Triassic palynology of New South Wales. Spec. Pub. Geol. Soc. Austral. 4, pp. 141 -155.
- INOSOVA K.J., KRUSINA A.M., SCHWARTZMAN E.G., 1976 - Atlas of the Upper Carboniferous and Lower Permian miospores and pollen from Donieck Basin. Moscow (in Russian).
- JACHOWICZ A., DYBOVA-JACHOWICZ S., 1987 - Carbiniferous and Permian palynology in Poland - a review. Prz. Geol. 5, Warsaw, pp. 259- 261.
- JACHOWICZ M., 1991 - Occurrence of Acritarcha in Cupper Shale in Poland. Micro-paleontology (in press).

- JANSONIUS J., 1962 - Palynology of Permian and Triassic Sediments of Peace River Area, Western Canada. *Palaeontographica B* 110/ 1-4, pp. 35-98.
- JEKHOWSKY B., GOUBIN N., 1964 - Subsurface palynology in Madagaskar: a stratigraphic sketch of the Permian, Triassic Jurassic of the Morondava Basin. In: A.T. Cross (Ed.): *Palynology in Oil Exploration. Spec. Pap. Soc. Econ. Palaeontologists and Mineralogists* 2, pp. 116-130.
- de JERSEY N.J., 1970 - Early Triassic miospores from the Revan Formation. *Publ. Geol. Surv. Queensland* 351, *Palaeont. Pap.* 19, 29 p.
- de JERSEY N.J., 1979 - Palinology of the Permian-Triassic transition in the Western Bowen Basin. *Publ. Geol. Surv. Queens.* 374.
- JURKIEWICZ H., 1962 - The foraminifera fauna from the Zechstein of Kajetanów, Tumlin and Gałżice. *Kwart. Geol.* 6/2, pp. 446-447 (in Polish).
- JURKIEWICZ H., 1966 - Foraminifera from the Upper Zechstein of Gałżice and Kajetanów. *Biull. IG* 195, pp. 159-194 (in Polish).
- KIUNTZEL M.K., 1965 - Palynological characteristic of the Upper Permian and Lower Triassic deposits in Weltuga River Basin and in the area between Wolga and Unzina Rivers in Kostromsk region. *Sbornik Stat. from Geology and Hydrogeology* 4, Moscow pp. 75-80 (in Russian).
- KLAUS W., 1955 - Über die Sporen diagnose des deutschen Zechstein salzes und des alpinen Salzgebirges. *Z. deutsch. geol. Ges.* 105/4, Hannover.
- KLAUS W., 1963 - Sporen aus dem südalpinen Perm. *Jb. Geol.* Bd. 106, pp. 229-363.
- KOTAŃSKA H., 1970 - A new genus of sporomorph from the Zechstein deposits of Poland. *Acta Palaeont. Pol.* 15/1, pp. 153-158.
- KOTAŃSKA H., KRASÓN J., 1966 - New sites of sporomorphs in the Zechstein deposits in Poland. *Bull. Acad. Pol. Sci., Ser. Sci. Geol. et Geogr.* 14/4, pp. 237-241.
- KOWALCZEWSKI Z., 1978 - Upper Permian deposits in northwestern part of the Góry Świętokrzyskie. In: R. Wagner, T. Piąkowski (Eds): *Guide to Excurs., Part 2., Symp. Central European Permian*, Warsaw pp. 33-40.
- KOWALCZEWSKI Z., RUP M., 1989 - Zechstein in the Holy Cross Mountains, Poland. *Biul. PIG* 362, pp. 5-39 (in Polish).
- KREMP E.M., BALME B.E., HELBY R.J., KYLE R.A., PLAYFORD T., PRICE P. L., 1977 - Carboniferous and Permian palynostratigraphy in Australia and Antarctica: a

- review. BMR Jour. of Austral. Geol. et Goephis. 2, pp. 177-208.
- KULETA M., 1985 - The petrographic-sedimentological investigation of the Buntsandstain in the NW margin of the Holy Cross Mts and Nida basin. State Geological Institute of Warsaw (unpublished) (in Polish).
- KULETA M., 1990 - Lower Triassic - Buntsandstain. In: Z. Kowalczewski (Ed.): The geological research of the Permian in the NW part of the Holy Cross Mts in the aspect of Cu, Pb, Zn, Fe and V occurrence. State Geological Institute of Warsaw (unpublished) (in Polish).
- KULETA M., RUP M., 1980 - The problem of the Permian/Triassic boundary in the NW part of the Holy Cross Mts, Poland. Kwart. Geol. 24/4, pp. 972 (in Polish).
- LESCHIK G., 1956 - Sporen aus dem Salzton des Zechsteins von Neuhof (bei Fulda). Palaeontographica B 100/4-6, pp. 122-142.
- LUBER A.A., WALTZ I.E., 1941 - Atlas of miospores and pollens of the Paleozoic USSR. Tr. VSEGEI 139, Leningrad (in Russian).
- MAHESHWARI H.K., BANERIJ J., 1975 - Lower Triassic palynomorphs from the Maritur Formation, West Bengal, India. Palaeontographica B 152/4-6, pp. 149-190.
- MASSARI F., CONTI M.A., FONTANA D., HELMOLD K., MARJOTTI N., NERI C., NICOSIA V., ORI G.G., PASINI M., PITFAN P., 1988 - The Val Gardena Sandstone and Bellerophon Formation in the Bletterbach Gorge (Alto Adige, Italy); biostratigraphy and sedimentology. Mem. Soc. Geol. Ital. 40, pp. 229-273.
- Mc GREGOR D.C., 1965 - Triassic, Jurassic and Lower Cretaceous spores and pollen of Arctic Canada. Geol. Surv. Canad. Pap. 66/65, 32 p.
- MEYEN S.V., 1987 - Fundaments of palaeobotany. Moscow, 403 p (in Russian).
- MOLIN W.A., 1989 - Palynostratigraphy of the Upper Permian on Kanin Peninsula. Palynol. taxon. in biostratigraphy, Moscow, pp. 135-140 (in Russian).
- MOLIN W.A., KOLODA N.A., 1972 - Upper Permian spore-pollen complexes from the northern part of the Russian Platform. Nauka, Moscow, 76 p (in Russian).
- ORŁOWSKA-ZWOLIŃSKA T., 1962 - A First Finding of Zechstein Sporomorphs in Poland. Kwart. Geol. 6/3, pp. 285-297 (in Polish).
- ORŁOWSKA-ZWOLIŃSKA T., 1984 - Palynostratigraphy of the Buntsandstein in sections of Western Poland. Acta Palaeont. Pol. 29/3-4, pp. 161-194.

- ORŁOWSKA-ZWOLIŃSKA T., 1985 - Palynological zones of the Polish Epicontinental Triassic., Bull. Pol. Acad. Sci., Earth Sci. 33/3-4, pp. 107-119.
- PATTISON J., SMITH D.B., WARRINGTON G., 1973 - A review of Late Permian and Early Triassic biostratigraphy in the British Islands. Can. Soc. Petrol. Geol. Mem. 2. In: A. Loghan, L.V. Hills (Eds.): The Permian and Triassic systems and their mutual boundary, pp. 220-260.
- PAWŁOWSKA K., 1964 - About the division of Zechstein in the Holy Cross Mts into the four sedimentary cycles. Prz. Geol. 9, pp. 367-371 (in Polish).
- PAWŁOWSKA K., 1978 - Zechstein in the Holy Cross Mts. In: R. Wagner T. Piątkowski (Eds.): Guide to Excurs., Part 2. Symp. Central European Permian, Warsaw, pp. 9-19.
- PIĘNKOWSKI G., 1987 - Permian/Triassic and Zechstein/Bundsandstein boundaries in Poland - a sedimentological approach. Abstr. and posters Inter. Symp. Zechstein '87, Bochum, pp. 42-43.
- PIĘNKOWSKI G., 1989 - The sedimentological criteria for the distinguishing of the Zechstein/Buntsandstein and Permian/ Triassic boundaries in Poland. Prz. Geol. 5, pp. 237-244 (in Polish).
- POTONIÉ R., 1956, 1958, 1960, 1966, 1970 - Synopsis der Gattungen der Sporae dispersae, Teil I-V. Beih. Geol. Jb. 23, 31, 39, 72, 87.
- POTONIÉ R., KLAUS W., 1954 - Einige Sporengattungen des alpinen Salzgebirges. Geol. Jb. 68, pp. 517-546.
- POTONIÉ R., KREMP G., 1954 - Die Gattungen der Palaeozoischen Sporen dispersae und ihre Stratigraphie. Geol. Jb. pp. 111- 194.
- POTONIÉ R., KREMP G., 1955, 1956a, 1956b - Die Sporae dispersae des Ruhrkarbon, Teil I-III. Palaeontographica B 98, pp. 1-136, 99, pp. 85-191, 100, pp. 65-121, Stuttgart.
- RICHTER-BERNBURG G., 1955 - Stratigraphische Gliederung des deutschen Zechstein. Z. Deutsch. Geol. Gess. 105.
- ROBINSON P.L., 1973 - Paleoclimatology and continental drift. In: D.H. Tarling, S.K. Runcorn (Eds.): Implications of Continental Drift to the Earth Sciences. Acad. Press. London/ New York, pp. 451-476.
- ROMANOWSKAYA G.M., 1963 - Spores and pollen. Palaeontology 2, Ural, Moscow (in Russian).

- ROMANOWSKAYA G.M., TABACHNIKOVA I.P., DRYAGINA L.L., 1973 - To the question on the Permian-Triassic boundary in Central Siberia (according to palynological data). Tr. VNIGRI 195, Leningrad, pp. 40-59 (in Russian).
- ROSPONDEK M., FIJAŁKOWSKA A., LEWANDOWSKA A., 1993 - The origin of the organic matter in Lower Silesian copper-bearing shales. Ann. Soc. Geol. Pol. 63, pp. 85-99.
- RUP M., 1985 - Correlation of the Upper Permian deposits in the Holy Cross region. State Geological Institute of Warsaw. (unpublished) (in Polish).
- SADOVNIKOV G.V., 1989 - About the Tutochansk horizont and volcanogenic Permian-Triassic beds in the Tungusk Basin. Tr. Inst. Geol. and Geophys. Akad. Nauk. SSSR 732, Moscow, pp. 313-319 (in Russian).
- SCHEURING B.W., 1970 - Palynologische und palynostratigraphische Untersuhungen des Keuper im Bölczentunnel. Schweiz. Palaont. Abh., Birkhauser Verlag, Basel.
- SENKOWICZOWA H., ŚLĄCZKA A., 1962 - Buntsandstein in the North margin of the Holy Cross Mountains. Roczn. Pol. Tow. Geol. 32/ 3, Cracow, pp. 313-331 (in Polish).
- SEVERCOVA I.A., 1966 - Spores and pollen from Permian deposits. In: Palaepalynology III, Nedra, Leningrad, pp. 114-134 (in Russian).
- SHU O., 1982 - Upper Permian and Lower Triassic palynomorphs from eastern Yunnan, China. Can. J. Earth Sci. 19/1, pp. 68-80.
- SHU O., 1986 - Palynology of Upper Permian and Lower Triassic strata of Fuyuan District, Eastern Yunnan. Acta Paleontol. Sinica 169, New S. A(9), pp. 103-122.
- SHU O., NORRIS G., 1988 - Spores and pollen from the Lower Triassic Heshauggou Formation, Shaanxi Province, Northern China. Rev. Palaeobot. and Palynol. 54, pp. 187-231.
- SMITH D.B., BRUNSTOM R.G.W., 1974 - Correlation of the Permian rock of the British Isles. Spec. Rep. Geol. Soc. 5, London.
- SMITH D.B., CROSBY X., 1979 - The regional and stratigrafical context of Zechstein 3 and 4 potash deposits in the southern North Sea and adjoinig areas. Econom. Geol. 74.
- SRIVASTAVA S.C., JHA N., 1988 - A Lower Triassic palynoassamblage from Budharam Area, Godavari Graben, Andhra Pradesh. Geophytology 18, pp. 124-125.
- TOTMAN-PARRIS J., CURTIS R.L., 1982 - Atmospheric Circulation, Upwelling and

- Organic-rich rocks in the Mesozoic and Cenozoic Eras. *Paleogeogr., Paleoclim., Paleoecol.* 40, pp. 31-66.
- UTTING J., 1979 - Pollen and spore assemblages from the Upper Permian of the North Luangwa Valley, Zambia. *Abstr. 4th Inter. Palynol. Congr.*, Lucknow 1976-1977, pp. 165-174.
- UTTING J., 1987 - Palynology of Uppermost (?) Carboniferous, Permian and Lower Triassic rocks of the Canadian Arctic Archipelago. *Abstr. and Pap. 11th Inter. Congr. Carbon. Stratigr. and Geol.*, Beijing 1987, pp. 384-386.
- VISSCHER H., 1966 - Palaeobotany of the Mezophytic III; Plant microfossils from the Upper Bunter of Hengelo, the Netherlands. *Acta Bot. Neerl.* 15, pp. 316-375.
- VISSCHER H., 1967 - Permian and Triassic palynology and concept of "Tethys twist". *Paleogeogr., Paleoccol., Paleoclim.* 3, pp. 151-168.
- VISSCHER H., 1971 - The Permian and Triassic of the Kingscourt outlier, Ireland. *Geol. Surv. of Ireland Spec. Pub.* 1.
- VISSCHER H., 1978 - Aspects of a palynological characterization of Late Permian and Early Triassic "standard" units of chronostratigraphical classification in Europe. *Proc. 4th Inter. Palyn. Confer.*, Lucknow 1976-1977, pp. 238-241.
- VISSCHER H., VAN DER ZWAN C.J., 1981 - Palynology of the Circum- Mediterranean Triassic: Phytogeographical and paleoclimatological implications. *Geol. Rund.* 70, pp. 625-634.
- VISSCHER H., BRUGMAN W.A., 1988 - The Permian-Triassic boundary in Southern Alps - a palynological approach. *Mem. Soc. Geol. Ital.* 34, pp. 121-128.
- WAGNER R., 1987 - Stratigraphy of the Uppermost Zechstein in Northern Poland. *Bull. Pol. Acad. Sci., Earth Sci.* 35/3, pp. 265-273.
- WAGNER R., PIĄTKOWSKI T.S., PERYT T.M., 1978 - The Polish Zechstein Basin. *Prz. Geol.* 12, Warsaw, pp. 673-686 (in Polish).
- WAGNER R., PIĄTKOWSKI T.S., PERYT T.M., 1981 - The evolution of the Zechstein sedimentary basin in Poland. *Proc. Inter. Symp. on Central European Permian*, Warszawa, pp. 59.
- WARIUKHINA L.M., 1971 - Spores and pollen of the bunter and coalbearing series of Permian and Triassic from north-east of the European part of USSR, Moscow (in Russian).

- ZBROJA St., 1990 - Lithostratigraphy of the Upper Permian. In: Z. Kowalczewski (Ed.):
The geological research of the Permian in the NW part of the Holy Cross Mts in the
aspect of Cu, Pb, Zn, Fe and V occurrence. State Geological Institute of Warsaw
(unpublished) (in Polish).
- ZHI W., JIAZHENG Z., 1987 - Carboniferous and Permian palynological assemblages in
Junggar Basin, Xinjiang. Abstr. 11th Inter. Congr. Carbon. Stratigr. and Sediment.,
Bijong 1987, pp. 387.

Appendix 1a. Range chart of all taxa in the Tumlin-Podgrodzie IG 1 borehole.

UPPER PERMIAN				LOWER TRIASSIC		LITHOSTRATIGRAPHY	
PZ1	PZ2	PZ3	Pzt	A0	A1	(after Zbroja, 1990 and Kuleta, 1990)	
T1 Ca1	A1d T1r	A1g	T3 T2r [Ca3]				
414.8	328.5	329.5	330.5	331.0	331.5	332.0	
DEPTH (in meters)							LITHOLOGY
							Localization of samples
++ *++		++ *+*					<i>Calamospora plicata</i>
++ *++		++					<i>Calamospora</i> sp.
+ *+ +		++					<i>Laevigatisporites</i> sp.
							<i>Punctatisporites triassicus</i>
							<i>Punctatisporites</i> sp.
							<i>Cyclotriletes triassicus</i>
							<i>Cyclotriletes microgranifer</i>
							<i>Verrucosisporites</i> sp.
							<i>Guttatisporites</i> sp.
							<i>Lycospora</i> cf. <i>permica</i>
							<i>Lycospora</i> sp.
							<i>Lundbladispora brevicula</i>
							<i>Lundbladispora</i> cf. <i>obsoleta</i>
							<i>Lundbladispora</i> sp.
							<i>Densoisporites palyfordii</i>
							<i>Densoisporites</i> sp.
							<i>Endosporites papillatus</i>
							<i>Laevigatosporites</i> sp.
							-- Sporites indet.
							<i>Florinites</i> sp.
							<i>Perisaccus granulatus</i>
							<i>Perisaccus</i> sp.
							<i>Potonieisporites simplex</i>
							<i>Potonieisporites</i> sp.
							<i>Cordaitina uralensis</i>
							<i>Cordaitina donetziana</i>
							<i>Cordaitina</i> sp.
							<i>Nuskoisporites dulhuntyi</i>
							<i>Nuskoisporites klausii</i>
							<i>Nuskoisporites</i> sp.
							<i>Trizonaesporites</i> sp.
							<i>Plicatipollenites indicus</i>
							<i>Protohaploxylinus samolovichii</i>
							<i>Protohaploxylinus jacobii</i>
							<i>Protohaploxylinus pantii</i>
*	* ++ *		++ **				<i>Protohaploxylinus</i> sp.
							<i>Strotersporites wilsoni</i>
							<i>Strotersporites richteri</i>
							<i>Strotersporites</i> sp.
							<i>Striatopodocarpites</i> sp.
							<i>Striatooabietites balmei</i>
							<i>Striatooabietites</i> sp.
*	* -- -		-- --				<i>Lueckisporites virkiae</i> NAA+AB
							<i>Lueckisporites virkiae</i> NAC
							<i>Lueckisporites virkiae</i> NBA+Bb
							<i>Lueckisporites virkiae</i> NBC
							<i>Lueckisporites virkiae</i> NC
							<i>Lueckisporites virkiae</i> f. with a "dark body"
*	* -- ** ***		-- -- **				<i>Lunatisporites noviaulensis</i>
							<i>Lunatisporites multiplex</i>
							<i>Lunatisporites labdacus</i>
							<i>Lunatisporites alatus</i>

	I	II	III	IV	SPORE-POLLEN ASSEMBLAGES
+	+	+	+	+	<i>Lunatisporites gracilis</i>
++	+	++	+	+	<i>Lunatisporites hexagonalis</i>
*	--	--	--	--	<i>Lunatisporites ortisei</i>
+	++	+			<i>Lunatisporites sp.</i>
+		+			<i>Vittatina vittifer</i>
+		+			<i>Vittatina hiltonensis</i>
+	++	++			<i>Vittatina sp.</i>
*	++	++			<i>Hamiapollenites sp.</i>
*	--	--			<i>Klausipollenites schaubergeri</i>
*	--	--			<i>Klausipollenites minimus</i>
*	**	**			<i>Klausipollenites sp.</i>
*	++	++			<i>Falcisporites zapfei</i>
+		+			<i>Paravesicaspora splendens</i>
+	**	**			<i>Platysaccus niger</i>
+	+	+			<i>Platysaccus papilionis</i>
+			--		<i>Platysaccus sp.</i>
*	++	++			<i>Illinites unicus</i>
+	++	++			<i>Illinites elegans</i>
+	++	++			<i>Illinites sp.</i>
+	++	++			<i>Jugasporites delasoucei</i>
+	--	--			<i>Jugasporites cf. parvus</i>
+	**	**			<i>Jugasporites paradelasoucei</i>
+	**	**			<i>Jugasporites sp.</i>
+				+	<i>Triadispora crassa</i>
+	**	**		**+	<i>Triadispora sp.</i>
+	--	--		--	<i>Limitisporites moersensis</i>
+	--	--			<i>Limitisporites leschiki</i>
+					<i>Limitisporites rectus</i>
+					<i>Limitisporites cf. parvus</i>
*	++	++			<i>Limitisporites sp.</i>
*	++	++			<i>Gardenasporites sp.</i>
+	++	++			<i>Cycadopites coxii</i>
+	++	++			<i>Cycadopites follicularis</i>
--	--	--			<i>Pollenites indet.</i>
-*					<i>Baltisphaeridium sp.</i>
**					<i>Micrhystridium sp.</i>
++					<i>Veryhachium sp.</i>

Lithology:		conglomerates		coarse sandstones		fine sandstones		mudstones, siltstones
		sandy mudstones		limestones		marly limestones		dolomites
		marly dolomites		carbonate nodules		mudstones clasts		sulphates

Palynology: — single positive sample, — more than one positive sample, -- single negative sample, -- more than one negative sample. Palynomorphs occurrence + 1-4 specimens, * 5-10 specimens, - 1-10%, = more than 10%.

Appendix 1b. Range chart of all taxa in the Jaworze IG 1 borehole.

---+**	*++	+-		<i>Illinites unicus</i>
+	+			<i>Illinites elegans</i>
+++	+	*		<i>Illinites sp.</i>
++	++		+ (row)	<i>Vitreisporites sp.</i>
--**--*	*++	+		<i>Jugasporites delasaucei</i>
+				<i>Jugasporites purus</i>
--*--*	++			<i>Jugasporites sp.</i>
+++++	++	++		<i>Triadispora sp.</i>
-----	+-	++		<i>Limitisporites moersensis</i>
+	+			<i>Limitisporites leschiki</i>
+	+			<i>Limitisporites rectus</i>
+	++			<i>Limitisporites cf. parvus</i>
++**	++			<i>Limitisporites sp.</i>
++	+++			<i>Gardenasporites sp.</i>
++	+			<i>Crustaeспорites sp.</i>
++				<i>Pakhapites sp.</i>
--*--*	*++	++	+ (row)	<i>Cycadopites coxii</i>
++	+	+		<i>Cycadopites follicularis</i>
++		+		* <i>Cycadopites sp.</i>
-----	---	--		- <i>Monosulcites sp.</i>
*****				- <i>Pollenites indet.</i>
****+*				+ <i>Baltisphaeridium debilispinum</i>
++ +				* <i>Baltisphaeridium sp.</i>
* ***				<i>Micrhystridium setasessitante</i>
*****+				+ <i>Micrhystridium recurvatum</i>
-----				* <i>Micrhystridium sp.</i>
-----				- <i>Veryhachium trispinoides</i>
-----				* <i>Veryhachium irregulare</i>
-***++				- <i>Veryhachium sp.</i>
				- <i>Acritarcha indet.</i>
				- "Thympanicysta"
	I		II	
	1		1	
				IV
				SPORE-POLLEN ASSEMBLAGES

For the legend see Appendix 1a.

Appendix 1c. Range chart of all taxa in the Zachełmie IG 1 borehole.

UPPER PERMIAN					LOWER TRIASSIC		LITHOSTRATIGRAPHY (after Zbroja, 1990 and Kuleta, 1990)	
PZ1	PZ2	T1r	T2r	Ca3	Pzt	A0	A1	DEPTH (in meters)
222	202	182	162	142	122	102	82	
222	202	182	162	142	122	102	82	
.....	LITHOLOGY
+	+	++			+			Localization of samples
+								<i>Calamospora</i> sp.
+								<i>Laevigatisporites</i> sp.
*								<i>Verrucosporites</i> sp.
+								<i>Lycospora</i> sp.
+								<i>Laevigatosporites</i> sp.
*	+	++			++			<i>Sporites</i> indet.
+	+	++			+			<i>Perisaccus granulatus</i>
+	+	++			++			<i>Perisaccus</i> sp.
+	+	++						<i>Potonieisporites simplex</i>
*	+	++						<i>Potonieisporites</i> sp.
+	+	++						<i>Cordaitina donetziana</i>
+	+	++						<i>Cordaitina</i> sp.
*	+	++						<i>Nuskoisporites dulhuntyi</i>
*	+	++						<i>Nuskoisporites klausii</i>
+	+	++						<i>Protohaploxylinus samoilovichii</i>
*	+	++						<i>Protohaploxylinus</i> sp.
+	+	++						<i>Strotersporites wilsoni</i>
+	+	++						<i>Strotersporites richteri</i>
+	+	++						<i>Striatoabietites balmei</i>
-	--							<i>Lueckisporites virkkiae</i> NAA+AB
+	+							<i>Lueckisporites virkkiae</i> NAC
+	++							<i>Lueckisporites virkkiae</i> NBa+Bb
+	++							<i>Lueckisporites virkkiae</i> NBC
+	++							<i>Lueckisporites virkkiae</i> NC
-	--				--			<i>Lueckisporites virkkiae</i> f. with a "dark body"
-	--				--			<i>Lunatisporites noviaulensis</i>
*	--				--			<i>Lunatisporites multiplex</i>
*	--				--			<i>Lunatisporites ortisei</i>
+	--				--			<i>Lunatisporites</i> cf. <i>alatus</i>
+	--				--			<i>Lunatisporites microsaccatus</i>
*	--				--			<i>Lunatisporites</i> sp.
+	--				--			<i>Vittatina vittifer</i>
+	--				--			<i>Hamiapollenites</i> sp.
+	--				--			<i>Klausipollenites schaubergeri</i>
+	--				--			<i>Falcisporites zapfei</i>
+	--				--			<i>Platysaccus niger</i>
+	--				--			<i>Platysaccus papilionis</i>
*	--				--			<i>Illinites unicus</i>
*	--				--			<i>Illinites elegans</i>
*	--				--			<i>Illinites</i> sp.
+	--				--			<i>Jugasporites delasaucei</i>
+	--				--			<i>Jugasporites</i> sp.
+	--				--			<i>Triadispora plicata</i>
+	--				--			<i>Triadispora</i> sp.
+	--				--			<i>Limitisporites moersensis</i>
+	--				--			<i>Limitisporites leschiki</i>
+	--				--			<i>Limitisporites</i> cf. <i>parvus</i>
+	--				--			<i>Limitisporites</i> sp.
+	--				--			<i>Gardenasporites leonardii</i>
+	--				--			<i>Gardenasporites</i> cf. <i>moroderi</i>
*	--				--			<i>Cycadopites</i> sp.
-	--				--			<i>Pollenites</i> indet.
	I				III			SPORE-POLLEN ASSEMBLAGES

For the legend see Appendix 1a.

Appendix 1d. Range chart of all taxa in the Zaczowice IG 1 borehole.

UPPER PERMIAN				LITHOSTRATIGRAPHY (after Zbroja, 1990)	
PZ1	PZ2	PZ3	Pzt	DEPTH (in meters)	
Ca1	T1r	T2r	T3 Ca3	LITHOLOGY	
					Localization of samples
+ +++					<i>Calamospora</i> sp.
++					<i>Laevigatisporites</i> sp.
+ + +		+ +			<i>Lycospora</i> sp.
++		+ ++			<i>Laevigatosporites</i> sp.
++++*++		+ ++			Sporites indet.
+**++ ++					<i>Potonieisporites simplex</i>
+ + +					<i>Potonieisporites</i> sp.
+ + + +					<i>Cordaitina uralensis</i>
++++*					<i>Cordaitina</i> sp.
-**--**+					<i>Nuskoisporites dulhuntyi</i>
+ + + +					<i>Nuskoisporites klausii</i>
++ *++*					<i>Nuskoisporites</i> sp.
++ + + ++					<i>Protohaploxylinus samoilovichii</i>
+++ +		+ ++			<i>Protohaploxylinus</i> sp.
		+ ++			<i>Strotersporites wilsoni</i>
		+ ++			<i>Strotersporites</i> sp.
-----*		*++*			<i>Lueckisporites virkkiae</i> NAa+Ab
+ +		+			<i>Lueckisporites virkkiae</i> NAc
++ *++*		+			<i>Lueckisporites virkkiae</i> NBa+Bb
+					<i>Lueckisporites virkkiae</i> NC
-----*		*	++		<i>Lunatisporites noviaulensis</i>
++*++*			+		<i>Lunatisporites labdacus</i>
+ * +**+			+ ++		<i>Lunatisporites</i> sp.
** +**. +					<i>Vittatina vittifer</i>
+ + ++					<i>Vittatina</i> sp.
+++++					<i>Hamiapollenites</i> sp.
-----*		----			<i>Klausipollenites schaubergeri</i>
+**++ +		++*			<i>Klausipollenites</i> sp.
++** +		++*			<i>Falcisporites</i> sp.
-**--**+		++ +			<i>Platysaccus niger</i>
+++ +		++			<i>Platysaccus</i> sp.
++ +					<i>Vitreisporites</i> sp.
-----		++*			<i>Jugasporites delasaucei</i>
++ * +		+			<i>Jugasporites</i> sp.
+ + +			+		<i>Triadispora</i> sp.
** + + + +		++*			<i>Limitisporites moersensis</i>
+ * +		+			<i>Limitisporites</i> sp.
+ + * +					<i>Gardenasporites</i> sp.
++*++ ++		++			<i>Cycadopites coxii</i>
-**--**		----			<i>Pollenites</i> indet.
*****					<i>Baltisphaeridium</i> sp.
-*- **					<i>Veryhachium</i> sp.
*****					<i>Acritarcha</i> indet.
I:		II:		SPORE-POLLEN ASSEMBLAGES	

b.c. - basal conglomerate

For the legend see Appendix 1a.

APPENDIX 2. Alphabetical listing of all miospore taxa identified

- Apiculatisporites apiculatus* f. *media* Dybova et Jachowicz 1957
Calamospora pedata Kosanke 1950
Calamospora plicata (Luber et Waltz 1941) Hart 1965
Chordasporites sp.
Cordaitina donetziana Inosova 1976
Cordaitina uralensis (Luber 1941) Dibner 1970
Crustaesporites cf. *latisulcatus* Lele et Maithy 1963
Cycadopites coxii Visscher 1966
Cycadopites follicularis Wilson et Webster 1946
Cyclotriletes microgranifer Mädler 1964
Cyclotriletes oligogranifer Mädler 1964
Densoisporites playfordii (Balme 1963) Dettmann 1963
Endosporites hexarecticulatus Klaus 1963
Falcisporites snopkove Visscher 1966
Falcisporites zapfei (Potonié et Klaus 1954) Leschik 1955
Florinites sp.
Gardenasporites heisseli Klaus 1963
Gardenasporites leonardii Klaus 1963
Gardenasporites cf. *moroperi* Klaus 1963
Gardenasporites cf. *oberrauchi* Klaus 1963
Gnetacaepollenites steevesi Jansonius 1962
Guttatisporites sp.
Hamiapollenites cf. *bifurcatus* Jansonius 1962
Illinites elegans Kosanke 1950
Illinites kosankei sp. nov.
Illinites unicus Kosanke 1950
Jugasporites delasaucei (Potonié et Klaus 1954) Leschik 1956
Jugasporites latus (Leschik 1956) Foster 1983
Jugasporites lueckoides Klaus 1963
Jugasporites paradelasaucei Klaus 1963
Jugasporites parvus (Klaus 1963) Foster 1983
Jugasporites purus (Leschik 1956) Tiwari et Singh 1984
Jugasporites schaubergeroides Klaus 1963
Klausipollenites decipiens Jansonius 1962
Klausipollenites minimus Góczán 1987
Klausipollenites staplinii Jansonius 1962
Laevigatisporites giganteus Dybova et Jachowicz 1957 f. *microsignus* f. nov.
Laevigatisporites minimalis f. *pulla* Dybova et Jachowicz 1957
Laevigatosporites vulgaris f. *minor* Loose 1934
Limitisporites moersensis (Grebe 1957) Klaus 1963
Limitisporites leschiki Klaus 1963
Limitisporites cf. *parvus* Klaus 1963
Limitisporites rectus Leschik 1956
Lueckisporites virkkiae Potonié et Klaus 1954
Lunatisporites acutus (Leschik 1956) Scheuring 1970
Lunatisporites alatus (Klaus 1963) comb. nov.

- Lunatisporites gracilis* (Jansonius 1962) comb. nov.
Lunatisporites hexagonalis (Jansonius 1962) Scheuring 1970
Lunatisporites labdacus (Klaus 1963) comb. nov.
Lunatisporites microsaccatus (Jansonius 1962) comb. nov.
Lunatisporites multiplex (Visscher 1966) Scheuring 1970
Lunatisporites noviaulensis (Leschik 1956) Scheuring 1970
Lunatisporites obex (Balme 1963) comb. nov.
Lunatisporites ortisei (Klaus 1963) Góczán 1987
Lunatisporites transversundatus (Jansonius 1962) comb. nov.
Lundbladispora brevicula Balme 1963
Lundbladispora cf. obsoleta Balme 1963
Lycospora permica (Inosova 1976) comb. nov.
Monosulcites sp.
Nuskoisporites dulhuntyi Potonié et Klaus 1954
Nuskoisporites klausii Grebe 1957
Pakhapites sp.
Paravesicaspora splendens (Leschik 1956) Klaus 1963
Perisaccus granulatus Klaus 1963
Pilaspora cf. *plurigenus* Balme et Henelly 1955
Platysaccus leschiki Hart 1960
Platysaccus niger Mädler 1964
Platysaccus papilionis Potonié et Klaus 1954
Plicatipollenites indicus Lele 1964
Potonieisporites simplex Wilson 1962
Protohaploxylinus jacobii (Jansonius 1962) Hart 1964
Protohaploxylinus cf. *latissimus* (Luber et Waltz 1941) Samoilovich 1953
Protohaploxylinus pantii (Jansonius 1962) Orłowska-Zwolińska 1984
Protohaploxylinus cf. *rhomboformis* (Polukhina 1960) Hart 1964
Protohaploxylinus samoilovichii (Jansonius 1962) Hart 1964
Protosacculina sp.
Punctatisporites triassicus Schulz 1964
Striatoabietites aytugii (Visscher 1966) Scheuring 1970
Striatoabietites balmei Klaus 1964
Striatopodocarpites sp.
Strotersporites richteri (Klaus 1955) Wilson 1962
Strotersporites wilsoni Klaus 1963
Triadispora crassa Klaus 1964 sensu Brugman 1979
Triadispora plicata Klaus 1964 sensu Brugman 1979
Triadispora visscheri (Visscher 1966) comb. nov.
Trizonaesporites grandis Leschik 1956
Verrucosisporites pseudomorulae Visscher 1966
Vesicaspora schemeli Klaus 1963
Vitreisporites sp.
Vittatina costabilis Wilson 1962
Vittatina hiltonensis Chaloner et Clarke 1962
Vittatina subsaccata Samoilovich 1953
Vittatina vittifera (Luber et Waltz 1941) Samoilovich 1953

APPENDIX 3. Taxonomy of selected taxa

1. *Cordaitina donetziana* Inosova, 1976 (Pl. I, fig. 19, 20).

DISCUSSION. Circular monosaccate pollen, 22-65 μm in size. Central body (c.b.) circular, saccus narrow on proximal side. Trilete mark usually invisible. C.b. exine thin, smooth or infrapunctate; saccus exine infrarecticulate.

REMARKS. Inosova has distinguished three forms (tenue, crassa and plicata) in this species but it is difficult to recognize them in Holy Cross material due to wors state of preservation.

AGE AND OCCURENCE. Upper Carboniferous - Lower Permian (common), Upper Permian (aundant) in: Poland and Ukraina.

2. *Cordaitina uralensis* (Luber, 1941) Dibner, 1970 (Pl. II, fig. 7).

DISCUSSION. Circular or oval monosaccate pollen, 50-80 μm in size. C.b. circular, saccus very narrow on proximal side. Saccus basis distinct. Tetrade mark invisible. C.b. exine smooth or finegranulate; saccus exine infrareticulate.

REMARKS. It differs from *C. donetziana* by the bigger size and luck of the tetrade mark.

AGE AND OCCURENCE. Middle Carboniferous - Lower Permian (common), Upper Permian (abundant) in: Poland, Ukraina, Russia (European and Asiatic parts) and Australia.

3. *Crustaeспорites globosus* Leschik, 1956 (Pl. VI, fig. 4).

DISCUSSION. Rounded trisaccate pollen, 48-75 μm in size. C.b. circular to slightly triangular; sacci small, reniform, may unite laterally. Tetrade mark invisible. C.b. exine divided into 6-8 irregular, smooth ribbs on proximal side; sacci exine reticulate.

AGE AND OCCURENCE. Upper Permian (common) in: Poland, Germany, India and Australia.

4. *Cyclotriletes microgranifer* Mädler, 1964 (Pl. I, fig. 3).

DISCUSSION. Circular spore, 52-58 μm in size. Trilete mark distinct, 2/3 spore radius lengh. Exine thick, granulate.

AGE AND OCCURENCE. Lower Triassic (abundant), Upper Triassic (rare) in: Poland, Germany and England.

5. *Cyclotriletes oligogranifer* Mädler, 1964 (Pl. I, fig. 7).

DISCUSSION. Circular spore, 72-81 μm in size. Trilete mark distinct, spore radius length. Exine thick, granulate with dense distribution of coniform grana.

REMARKS. It differs from *C. microgranulatus* by the bigger size and coarse ornamentation.

AGE AND OCCURENCE. Lower Triassic (abundant), Upper Triassic (rare) in: Poland, Germany, England and Ukraina.

6. *Densoisporites playfordii* (Balme, 1963) Dettmann, 1963 (Pl. I, fig. 17).

DISCUSSION. Circular to slightly triangular, cavate spore, 40-65 μm in size. Wide cingulum in equatorial zone. Trilete mark distinct, extending to inner spore margin. Exine rough, thickened in equatorial margin.

AGE AND OCCURENCE. Upper Permian (rare), Lower Triassic (abundant) in: Poland, Eastern Greenland, Canadian Arctic Archipelago, Russia (Europaen and Asiatic part), Pakistan, India, Israel, Madagaskar, Australia and Tasmania.

7. *Falcisporites zapfei* (Leschik, 1956) Klaus, 1963 (Pl. IV, fig. 16).

DISCUSSION. Haploxytonoidal bisaccate pollen, 30-60 μm in size. C.b. circular, sacci less than hemispherical in shape, with distal inclination. Sacci basis distinct. Narrow, long sulcus on distal side. C.b. exine reticulate more clear on proximal side; sacci exine irregular, reticulate.

AGE AND OCCURENCE. Upper Permian (common), Lower Triassic (rare) in: Europe and Canadian Arctic Archipelago.

8. *Gardenasporites heisseli* Klaus, 1963 (Pl. VI, fig. 3).

DISCUSSION. Slight diploxytonoid bisaccate pollen, 40-80 μm in size. C.b. longitudinally oval, sacci hemispherical with distal inclination. Sacci basis convex. Long (40 μm), monolete mark on proximal pole. C.b. exine thin, reticulate; sacci exine finereticulate.

AGE AND OCCURENCE. Upper Permian (common), Lower Triassic (sporadic) in: Poland, Germany, North Sea Basin, Hungary and Italy.

9. *Illinites unicus* Kosanke, 1950 (Pl. V, fig. 6).

DISCUSSION. Haploxytonoid bisaccate pollen, 32-76 μm in size. C.b. circular or

longitudinally oval. Sacci hemispherical or less then hemispherical with distal inclination. Sacci basis convex or straight, parallel. Small, trilete mark with shorter one arm. C.b. exine fine-granulate, reduced in round mark area; sacci exine reticulate, built from big, sometimes irregular meshes.

AGE AND OCCURENCE. Upper Carboniferous (rare), Upper Permian (abundant), Lower Triassic (sporadic) in: Poland, Germany, England, Italy and USA.

10. *Jugasporites delasaucei* (Potonié et Klaus, 1954) Leschik, 1956 (Pl. V, fig. 4).

DISCUSSION. Diploxylonoid bisaccate pollen, 30-65 μm in size. C.b. transversaly oval, sacci hemispherical, bigger then c.b. Distal zone narrow. Slight convex sacci basis present. Tetrade mark in shape of dilet or trilet with one shorter arm on proximal pole. C.b. exine thick, infrapapillate on proximal side and smooth - on distal one. There is area without exoexine round tetrade mark. Sacci exine fine, reticulate.

AGE AND OCCURENCE. Upper Permian (rare), Upper Permian (abundant), Lower Triassic (sporadic) in: Poland, Western Europa, Italy, Ukraina and China.

11. *Jugasporites lueckoides* Klaus 1963 (Pl. V, fig. 9).

DISCUSSION. Haploxylonoid bisaccate pollen, 30-50 μm in size. C.b. transversaly oval. Sacci small, semilunar in shape. Distal zone widh. Monolete or diletete tetrade mark, 1/3 c.b. radius lengh. C.b. exine infrapapillate or infrarticulate. Area round tetrade mark without exoexine. Sacci exine reticulate.

REMARKS. It differs from other *Jugasporites* species by big, oval area roud tetrade mark, which conforms it to *Lueckisporites*.

AGE AND OCCURENCE. Upper Permian (not numerous) in: Poland, England and Italy.

12. *Jugasporites paradelasaucei* Klaus 1963 (Pl. V, fig. 15).

DISCUSSION. Diploxylonoid bisaccate pollen, 35-82 μm in size. C.b. circular. Sacci less then hemispherical in shape. Distal zone wide. Straight sacci basis present. Diletete or trilete, with one shorter arm, tetrade mark surrounded with area of reduced exoexine. C.b. exine infrapappillate to reticulate; sacci exine reticulate.

REMARKS. It differs from *J. delasaucei* by more circular shape of c.b. and wider distal zone.

AGE AND OCCURENCE. Upper Permian (abundant) in: Poland, England and Italy.

13. *Jugasporites schaubergeroides* Klaus 1963 (Pl. V, fig. 13).

DISCUSSION. Haploxylonoid bisaccate pollen, 26-48 μm in size. C.b. circular or oval. Sacci smaller than c.b., falciform in shape. Convex sacci basis distinct. Distal area of 1/2 c.b. radius width. Trilete tetrade mark with arms 9-11 μm long surrounded with small area of reduced exine. C.b. exine granulate on proximal side and smooth on distal one. Sacci exine reticulate.

REMARKS. It differs from the other *Jugasporites* species by small size and falciform sacci that conforms it to *Klausipollenites schaubergeri*.

AGE AND OCCURENCE. Upper Permian (not numerous) in: Poland, England, Hungary and Italy.

14. *Klausipollenites schaubergeri* (Potonié et Klaus 1954) Jansonius 1962 (Pl. IV, fig. 8).

DISCCUSION. Haploxylonoidd bisaccate pollen, 20-61 μm in size. C.b. circular or oval. Sacci small, less than hemispherical to semilunar in shape. Sometimes sacci basis united in equatorial zone. Tetrade mark invisible. C.b. exine infragranulate on proximal side and infrareticulate on distal one, conform to sacci exine.

AGE AND OCCURENCE. Lower Permian (regular), Upper Permian (adundant), Lower Triassic (frequent) in: Poland, Western and Southern Europe, Russia (European part), Ukraina, India, Central and Southern Africa and Australia.

15. *Limitisporites moersensis* (Grebe 1957) Klaus 1963 (Pl. V, fig. 18).

DISCUSSION. Haploxylonoid bisaccate pollen, 40-60 μm in size. C.b. circular or transversaly oval. Sacci hemispherical, with distal inclination. Straight or curved monolete mark 3/4 c.b. radius lengh with narrow sutura on proximal pole. C.b. exine fine-granulate to irregular, reticulate. Sacci exine infrareticulate.

AGE AND OCCURENCE. Upper Permian (abundant) in: Poland, Western Europe and Italy.

16. *Limitisporites rectus* Leschik 1956 (Pl. V, fig. 21).

DISCUSSION. Haploxulonoid bisaccate pollen, 25-66 μm in size. C.b. circular or transveraly oval. Sacci hemispherical to semilunar, with distal inclination. Straight or fusiform sacci basis

present. Distinct, straight monolete mark of different length on proximal pole. C.b. exine fine-punctate to granulate; sacci exine irregular, reticulate.

REMARKS. Comparatively short, straight tetrade mark and characteristic sacci basis distinguish it from other *Limitisporites* species.

AGE AND OCCURENCE. Lower Permian (regular), Upper Permian (frequent) in: Poland, Western Europa, Hungary, Italy, Ukraina, Russia (European part), Tanzania, Australia and Brazil.

17. *Lueckisporites virkkiae* Potonié et Klaus 1954 (Pl. VII, fig. 1-9).

DISCUSSION. Diploxytonoid bisaccate pollen, 40-95 μm in size. C.b. circular or transversaly oval. Sacci less then hemispherical. Sacci basis straight or slightly convex. Distal zone comparatively wide. Monolete tetrade mark 1/3-1/2 c.b. radius length on proximal pole. C.b. exine divided in two hemispherical taeniae, with infrareticular ornamentation. Sacci exine reticulate.

REMARKS. Visscher (1971) introduced the "palynodeme" idea and applied it to *Lueckisporites* genus. He distinguished, on the basis of changes in the exoexine structure and the shape of sacci, several palynological norms. Only four of them (A, B, C and E) was recognized in the Holy Cross Mts. It shoud be mentioned that the "palynodeme" concept has been modified last time.

AGE AND OCCURENCE. Lower Permian (sporadic), Upper Permian (abundant), Lower Triassic (sporadic) in: Poland, Western and Southern Europe, Hungary, Russia (European part), China, Pakistan, Madagaskar, Central and Southern Africa.

18. *Lunatisporites noviaulensis* (Leschik 1956) Scheuring 1970 (Pl. IV, fig. 1).

DISCUSSION. Diploxytonoid bisaccate pollen, 60-120 μm in size. C.b. circular or transversaly oval. Sacci hemispherical or bigger in shape. Sacci bassis not always visible. Distal zone broad. Monolete mark on proximal pole. C.b. exine divided into four taeniae: two polar and two equatorial. Surface of taeniae smooth or infrareticulate. Sacci exine reticulate.

AGE AND OCCURENCE. Upper Permian-Lower Triassic (abundant), Middle Triassic (frequent), Upper Triassic (rare) in: Poland, Western and Southern Europe, Hungary, Russia (European part), Ukraina, China, India, Pakistan, Madagaskar, Central Africa and Australia.

19. *Lundbladispora brevicula* Balme 1963 (Pl. I, fig. 13).

DISCUSSION. Cavate, rounded-triangular spore, 32-40 μm in size. Distinct trilete mark accompanied by broad, sometimes undating lips. Exine thick, covered with conical elements, distributed over distal and in marginal part of proximal side.

AGE AND OCCURENCE. Lower Triassic (common) in: Poland, Pakistan and Australia.

20. *Nuskoisporites dulhuntyi* Potonié et Klaus 1954 (Pl. II, fig. 4).

DISCUSSION. Circular to oval monosaccate pollen, 50-200 μm in size. C.b. circular. Saccus width of 1/3-1/2 c.b. radius. Trilete mark with short arms sometimes surrounded with darker, circular area. C.b. exine reticulate, built from big, polygonal meshes. Saccus exina reticulate, radially elongated.

AGE AND OCCURENCE. Upper Permian (common), Lower Traissic (rare) in: Poland, Western and Southern Europe, Hungary, Russia (European part) and China.

21. *Nuskoisporites klausii* Grebe 1957 (Pl. II, fig. 5).

DISCUSSION. Big, circular monosaccate pollen, 150-280 μm in size. C.b. small, circular. Trilete mark with arms 1/2 c.b. radius length surrounded by daker area. Exine of c.b. and saccus similar reticulate to infragranulate.

AGE AND OCCURENCE. Upper Permian (common) in: Poland, Germany and England.

22. *Paravesicaspora splendens* Klaus 1963 (Pl. IV, fig. 20).

DISCUSSION. Haploxylonoid bisaccate pollen, 65-89 μm in size. C.b. rhomboidal, longitudinally elongated. Sacci hemispherical, with distal inclination. Their basis may unite on distal side. Long, straight, distal sulcus. C.b. exine infrareticulate to infrabaculate on proximal side and fine-infrareticulate on dsital one. Sacci exine reticulate.

AGE AND OCCURENCE. Lower Permian (rare), Upper Permian (frequent) in Poland, Western Europa, Italy and Zair.

23. *Perisaccus granulatus* Klaus 1963 (Pl. I, fig. 16).

DISCUSSION. Circular to oval monosaccate pollen, 40-85 μm in size. C.b. usually circular. Saccus width differentiated from 2/3 c.b radius to 2/3 c.b. diameter. Trilete mark with arms of unequal lengh. C.b. exine fine-punctate to infrareticulate; saccus exine granulate.

AGE AND OCCURENCE. Upper Permian (common) in: Poland, Western Europe and Italy.

24. *Protohaploxylinus pantii* (Jansonius 1962) Orłowska-Zwolińska 1984 (Pl. III, fig. 6).
 DISCUSSION. Diploxylonoidal bisaccate pollen, 65-82 μm in size. C.b. circular or oval. Sacci semispherical with distal inclination. Distinct sacci basis. Some specimens have undistinct monolete mark. C.b. exine separated into 6-9 (usually 7) taeniae. Surface of taeniae smooth or infragranulate. Sacci exine infrareticulate.

AGE AND OCCURENCE. Upper Permian (rare), Lower Triassic (abundant), Middle Triassic (rare) in: Poland, England, Italy, Western Canada, India, Pakistan, Madagaskar and Tanzania.

25. *Protohaploxylinus samoilovichii* (Jansonius 1962) Hart 1964 (Pl. III, fig. 1).

DISCUSSION. Diploxylonoid bisaccate pollen, 40-95 μm in size. C.b. circular or longitudinally oval. sacci hemispherical or less then hemispherical in shape, with distal inclination. Distal zone narrow. Distinct sacci basis. Monolete mark 2/3 c.b. radius length on proximal pole. Exine on proximal side of c.b. divided into 8- 13, rather broad (3-7 μm) taeniae separated by narrow striae. Taeniae are regular, parallel to one another over whole c.b. Sacci exine infrareticulate, folded in vicinity of sacci attachments.

REMARKS. It is distinguished by the continuos parallel taeniae and distinct sacci basis.

AGE AND OCCURENCE. Lower Permian (rare), Upper Permian (common), Lower Triassic (abundant), Middle Triassic (frequent) in: Poland, Western Europa, Italy, Eastern Greenland, Western Canada, Tanzania, Australia and Tasmania.

26. *Punctatisporites triassicus* Schulz 1964 (Pl. I, fig. 8).

DISCUSSION. Circular spore, 56-60 μm in size. Distinct trilete mark with arms 2/3 spore radius length, sometimes divided at ends. Exine thick with fine, punctate ornamentation.

AGE AND OCCURENCE. Lower Triassic (abundant) in: Poland, Germany, Ukraina and Romania.

27. *Striatoabietites balmei* Klaus 1964 (Pl. III, fig. 7).

DISCUSSION. Diploxylonoid to slightly haploxylonoid bisaccate pollen, 45-72 μm in size. C.b. oval. Sacci hemispherical. Distal zone broad. Small, trilete or dilete mark usually

invisible. C.b. exine divided to numerous, narrow ribbs with smooth surface which continue over c.b. Sacci exine fine, infrareticulate.

AGE AND OCCURENCE. Upper Permian (not frequent), Lower-Middle Triassic (abundant), Upper Triassic (rare) in: Poland, Western Europe, Italy, Western Canada, Russia (Ural), Australia.

28. *Strotersporites richteri* (Klaus 1955) Wilson 1962 (Pl. III, fig. 8).

DISCUSSION. Diploxylonoid bisaccate pollen, 52-80 μm in size. C.b circular or oval. sacci less then hemispherical in shape. Straight or slight curved monolete mark on proximal pole. C.b. exine divided into numerous, discontinous ribbs with distinc, transverse segmentation. Sacci exine reticulate built from big, polygonal meshes.

REMARKS. It is distinguished by the presence of the transverse segmantation of ribbs exine.

AGE AND OCCURENCE. Upper Permian (common), Lower Triassic (frequent) in: Poland, Western Europe, Hungary, Italy, Eastern Greenland, Western Canada, Canadian Arctic Archipelago, USA.

29. *Strotersporites wilsoni* Klaus 1963 (Pl. III, fig. 11).

DISCUSSION. Slight diploxylonoid bisaccate pollen, 60-115 μm in size. C.b. subcircular. Sacci semilunar, with distal inclination. Distal zone narrow. Monolete mark 1/3 c.b. radius lengh on proximal pole. C.b. exine divided into numerous, narrow ribbs with granulate surface. Sacci exine reticulate built from big, distinct, polygonal meshes.

AGE AND OCCURENCE. Upper Permian (common) in: Poland, England and Italy.

30. *Vittatina vittifera* (Luber et Waltz 1941) Samoilovich 1953 (Pl. IV, fig. 11).

DISCUSSION. Transversaly oval pollen with reduced or without sacci, 40-65 μm in size. Exine of c.b. divided into 12-14 narrow ribbs with smooth surface.

AGE AND OCCURENCE. Lower Permian (common, locally abundant), Upper Permian (frequent), Lower Triassic (sporadic) in: Poland, Russia (European part and Eastern Ural), Italy and Tanzania.

Explanation to the Plates

All specimens x 500 if no other magnification is given.

PLATE I

- Fig. 1. *Calamospora plicata* (Luber et Waltz) Hart;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 2. *Calamospora pedata* Kosanke;
the Siodła IG 1 borehole, 212,0 m; Zechstein, PZ1 (T1r).
- Fig. 3. *Cyclotriletes microgranifer* Mädler;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).
- Fig. 4. *Laevigatisporites minimalis* f. *pulla* Dybova et Jachowicz;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 5. *Laevigatisporites giganteus* Dybova et Jachowicz f. *microsignus* f. nov.;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.
- Fig. 6. *Verrucosisporites pseudomorulae* Visscher;
the Siodła IG 1 borehole, 212,0 m; Zechstein, PZ1 (T1r).
- Fig. 7. *Cyclotriletes oligogranifer* Mädler;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).
- Fig. 8. *Punctatisporites triassicus* Schulz;
the Tumlin-Podgrodzie IG 1 borehole, 190,7 m; Buntsandstein (complex A1).
- Fig. 9. *Apiculatisporites apiculatus* f. *media* Dybova et Jachowicz;
the Stachura IG 1 borehole, 657,3 m; Zechstein, PZ1 (A1g).
- Fig. 10. *Florinites* sp.;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).
- Fig. 11. *Laevigatosporites vulgaris* f. *minor* Loose;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.
- Fig. 12. *Lycospora permica* (Inosova) comb. nov.;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 13. *Lundbladispora brevicula* Balme;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).

Fig. 14. *Lundbladispora* cf. *obsoleta* Balme;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).

Fig. 15. *Guttatisporites* sp.;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).

Fig. 16. *Perisaccus granulatus* Klaus;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).

Fig. 17. *Densoisporites playfordii* (Balme) Dettmann;
the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1).

Fig. 18. *Potonieisporites simplex* Wilson;
the Radwanów IG 1 borehole, 1630,5 m; Zechstein, PZ1 (Ca1).

Fig. 19. *Endosporites hexarecticulatus* Klaus;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).

Fig. 20, 21. *Cordaitina donetziana* Inosova;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).

PLATE II

Fig. 1, 3. *Trizonaesporites grandis* Leschik;
the Jaworze IG 1 borehole, 369,5 m; Zechstein, PZ1 (A1g) (Fig. 1. - x 1000).

Fig. 2. *Plicatipollenites indicus* Lele;
the Jaworze IG 1 borehole, 369,5 m; Zechstein, PZ1 (A1g).

Fig. 4. *Nuskoisporites dulhuntyi* Potonié et Klaus;
the Łączna-Zaszosie IG 1 borehole, 435,2 m; Zechstein, PZ1 (T1r) (x 750).

Fig. 5. *Nuskoisporites klausii* Grebe;
the Łączna-Zaszosie IG 1 borehole, 435,2 m; Zechstein, PZ1 (T1r) (x 1000).

Fig. 6. *Crucisaccites* cf. *latisulcatus* Lele et Maithy;
the Łopuszno IG 1 borehole, 1761,9 m; Zechstein, PZ1 (Ca1).

Fig. 7. *Cordaitina uralensis* (Luber) Dibner;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).

PLATE III

- Fig. 1. *Protohaploxylinus samoilovichii* (Jansonius) Hart;
the Stachura IG 1 borehole, 665,6 m; Zechstein, PZ1 (A1g).
- Fig. 2. *Protohaploxylinus cf. latissimus* (Luber et Waltz) Samoilovich;
the Łaczna-Zaszosie IG 1 borehole, 435,2 m; Zechstein, PZ1, (T1r) (x 750).
- Fig. 3. *Protohaploxylinus jacobii* (Jansonius) Hart;
the Stachura IG 1 borehole, 665,6 m; Zechstein, PZ1 (A1g) (x 750).
- Fig. 4. *Protohaploxylinus cf. rhomboeformis* (Polukhina) Hart;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).
- Fig. 5. *Striatoabietites aytugii* (Visscher) Scheuring;
the Łaczna-Zaszosie IG 1 borehole, 432,5 m; Zechstein, PZ1 (T1r).
- Fig. 6. *Protohaploxylinus pantii* (Jansonius) Orłowska-Zwolińska;
the Łaczna-Zaszosie IG 1 borehole, 432,5 m; Zechstein, PZ1 (T1r).
- Fig. 7. *Striatoabietites balmei* Klaus;
the Stachura IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 8. *Strotersporites richteri* (Klaus) Wilson;
the Stachura IG 1 borehole, 665,6 m; Zechstein, PZ1 (A1g).
- Fig. 9. *Lunatisporites alatus* (Klaus) comb. nov.;
the Jaworze IG 1 borehole, 369,5 m; Zechstein, PZ1 (A1g).
- Fig. 10. *Lunatisporites gracilis* (Jansonius) comb. nov.;
the Łaczna-Zaszosie IG 1 borehole, 432,5 m; Zechstein, PZ1 (T1r).
- Fig. 11. *Strotersporites wilsoni* Klaus;
the Siodła IG 1 borehole, 183,2 m; Zechstein, Pzt.
- Fig. 12. *Lunatisporites obex* (Balme) comb. nov.;
the Podgace IG 1 borehole, 97,7 m; Zechstein, PZ1 (T1r).
- Fig. 13. *Striatopodocarpites* sp.;
the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1).
- Fig. 14. *Lunatisporites acutus* Leschik;
the Łaczna-Zaszosie IG 1 borehole, 432,5 m; Zechstein, PZ1 (T1r).
- Fig. 15. *Lunatisporites hexagonalis* (Jansonius) Scheuring;
the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1).

- Fig. 16. *Lunatisporites labdacus* (Klaus) comb. nov.;
the Radwanów IG 1 borehole, 1630,9 m; Zechstein, PZ1 (Ca1).
- Fig. 17. *Lunatisporites microsaccatus* (Jansonius) comb. nov.;
the Zachełmie IG 1 borehole, 185,6 m; Zechstein, PZ1 (T1r).
- Fig. 18. *Lunatisporites ortisei* (Klaus) Góczán;
the Stachura IG 1 borehole, 657,9 m; Zechstein, PZ1 (A1g).

PLATE IV

- Fig. 1. *Lunatisporites noviaulensis* (Leschik) Scheuring;
the Łopuszno IG 1 borehole, 1761,8 m; Zechstein, PZ1 (Ca1).
- Fig 2, 6. *Vittatina subsaccata* Samoilovich;
Fig 2. - the Radwanów IG borehole, 1645,7 m; Zechstein, PZ1 (Ca1); Fig. 6. - the Stachura
IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 3. *Vittatina costabilis* Wilson;
the Siodła IG 1 borehole, 182,5 m; Zechstein, PZt.
- Fig. 4. *Lunatisporites transversundatus* (Jansonius) comb. nov.;
the Łopuszno IG 1 borehole, 1761,9 m; Zechstein, PZ1 (Ca1).
- Fig. 5. *Vittatina hiltonensis* Chaloner et Clarke;
the Łopuszno IG 1 borehole, 1735,5 m; Zechstein, PZ1 (Ca1).
- Fig. 7. *Lunatisporites multiplex* (Scheuring) comb. nov.;
the Łopuszno IG 1 borehole, 1744,8 m; zechstein, PZ1 (Ca 1).
- Fig. 8. *Klausipollenites schaubergeri* (Potonié et Klaus) Jansonius;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).
- Fig. 9, 10. *Klausipollenites minimus* Góczán;
the Podgace IG 1 borehole, 97,7 m; Zechstein, PZ1 (T1r);
Fig. 9. - the distal side, Fig. 10. - the proximal side.
- Fig. 11. *Vittatina vittifera* (Luber et Waltz) Samoilovich;
the Radwanów IG 1 borehole, 1645,7 m; Zechstein, PZ1 (Ca1).
- Fig. 12. *Hamiapollenites cf. bifurcatus* Jansonius;
the Tumlin-Węgle IG 1 borehole, 125,7 m; Zechstein, PZ1 (Ca1).

- Fig. 13. *Klausipollenites staplinii* Jansonius;
the Stachura IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 14. *Protosacculina* sp.;
the Stachura IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 15. *Klausipollenites decipiens* Jansonius;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1) (x 1000).
- Fig. 16. *Falcisporites zapfei* (Potonié et Klaus) Leschik;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.
- Fig. 17. *Vesicaspora schemeli* Klaus;
the Stachura IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 18, 19. *Falcisporites snopkove* Visscher;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1);
Fig. 18. - the distal side, Fig. 19. - the proximal side.
- Fig. 20. *Paravesicaspora splendens* (Leschik) Klaus;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt (x 750).

PLATE V

- Fig. 1. *Platysaccus leschiki* Hart;
the Stachura IG 1 borehole, 602,5 m; Zechstein, PZ3 (Ca3) (x 750).
- Fig. 2. *Platysaccus papilionis* Potonié et Klaus;
the Jaworze IG 1 borehole, 369,5 m; Zechstein, PZ1 (A1g).
- Fig. 3. *Platysaccus niger* Mädler;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 4. *Jugasporites delasaucei* (Potonié et Klaus) Leschik;
the Łopuszno IG 1 borehole, 1761,5 m; Zechstein, PZ1 (Ca1).
- Fig. 5. *Jugasporites parvus* (Klaus) Foster;
the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1).
- Fig. 6. *Illinites unicus* Kosanke;
the Siodła IG 1 borehole, 182,3 m; Zechstein, Pzt (x 750).
- Fig. 7. *Jugasporites latus* (Leschik) Foster;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).

- Fig. 8. *Illinites elegans* Kosanke;
the Stachura IG 1 borehole, 665,4 m; Zechstein, PZ1 (A1g).
- Fig. 9. *Jugasporites lueckoides* Klaus;
the Siodła IG 1 borehole, 183,2 m; Zechstein, Pzt.
- Fig. 10. *Illinites kosankei* sp. nov. (holotype);
the Siodła IG 1 borehole, 183,2 m; Zechstein, Pzt.
- Fig. 11. *Jugasporites purus* (Leschik) Tiwari et Singh;
the Stachura IG 1 borehole, 665,5 m; Zechstein, PZ1 (T1r).
- Fig. 12. *Vitreisporites* sp.;
the Stachura IG 1 borehole, 639,7 m; Zechstein, PZ1 (A1g).
- Fig. 13. *Jugasporites schaubergeroides* Klaus;
the Siodła IG 1 borehole, 183,2 m; Zechstein, Pzt.
- Fig. 14. *Triadispora visscheri* (Visscher) comb. nov.;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 15. *Jugasporites paradelasaucei* Klaus;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.
- Fig. 16. *Triadispora plicata* Klaus;
the Łopuszno IG 1 borehole, 1761,9 m; Zechstein, PZ1 (Ca1).
- Fig. 17, 19. *Triadispora crassa* Klaus;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).
- Fig. 18. *Limitisporites moersensis* (Grebe) Klaus;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1) (x 750).
- Fig. 20. *Limitisporites leschiki* Klaus;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).
- Fig. 21. *Limitisporites rectus* Leschik;
the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1).
- Fig. 22. *Gardenasporites* cf. *moroderi* Klaus;
the Zachełmie IG 1 borehole, 185,6 m; Zechstein, PZ1 (T1r).
- Fig. 23. *Gardenasporites* cf. *oberrauchi* Klaus;
the Siodła IG 1 borehole, 183,2 m; Zechstein, Pzt (x 750).

PLATE VI

- Fig. 1. *Gardenasporites leonardii* Klaus;
the Siodła IG 1 borehole, 212,2 m; Zechstein, PZ1 (T1r) (x 750).
- Fig. 2. *Limitisporites cf. parvus* Klaus;
the Zachełmie IG 1 borehole, 185,6 m; Zechstein, PZ1 (T1r).
- Fig. 3. *Gardenasporites heisseli* Klaus;
the Siodła IG 1 borehole, 212,2 m; Zechstein, PZ1 (T1r).
- Fig. 4. *Crustaeспорites globosus* Leschik;
the Łączna-Zaszosie IG 1 borehole, 435,7 m; Zechstein, PZ1 (T1r).
- Fig. 5. *Pakhapites* sp.;
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1).
- Fig. 6. *Gnetaceaepollenites steevesi* Jansonius;
the Ostojów IG 1 borehole, 262,0 m; Buntsandstein (complex A1).
- Fig. 7. *Cycadopites coxii* Visscher;
the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.
- Fig. 8. *Cycadopites follicularis* Wilson et Webster;
the Radwanów IG 1 borehole, 1645,0 m; Zechstein, PZ1 (Ca1).
- Fig. 9. *Monosulcites* sp.;
the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1).
- Fig. 10, 22, 25. *Michrystridium cf. stellatum* Deflandre;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1) (x 750).
- Fig. 11. *Leiosphaeridia* sp.;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).
- Fig. 12. *Michrystridium recurvatum* Valensi;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).
- Fig. 13. *Chordasporites* sp.;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 14. *Michrystridium setasessitante* Jansonius;
the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).
- Fig. 15. *Baltisphaeridium debilispinum* Wall et Downie;
the Radwanów IG 1 borehole, 1631,5 m; Zechstein, PZ1 (Ca1).

Fig. 16. *Baltisphaeridium* sp.;

the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1) (x 750).

Fig. 17-19. *Veryhachium irregulare* Jekhowsky;

the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).

Fig. 20. *Veryhachium* sp.;

the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1).

Fig. 21. *Veryhachium trispinoides* (Jekhowsky) comb. nov.;

the Radwanów IG borehole, 1630,0 m; Zechstein, PZ1 (Ca1) (x 750).

Fig. 23. *Baltisphaeridium* sp.;

the Radwanów IG 1 borehole, 1647,0 m; Zechstein, PZ1 (Ca1) (x 750).

Fig. 24. *Pilaspora* cf. *plurigenus* Balme et Hennelly;

the Siodła IG 1 borehole, 182,5 m; Zechstein, Pzt.

Fig. 26. *Michrystridium* sp.;

the Łączna-Zaszosie IG 1 borehole, 329,7 m; Buntsandstein (complex A1) (x 1000).

Fig. 27. *Baltisphaeridium longispinosum* (Eisenack) Eisenack;

the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1) (x 1000).

PLATE VII

Fig. 1. *Lueckisporites virkkiae* Potonié et Klaus norm (N) Aa (after Visscher);

the Radwanów IG 1 borehole, 1630,0 m; Zechstein, PZ1 (Ca1) (x 750).

Fig. 2. *Lueckisporites virkkiae* Potonié et Klaus NAb;

the Łopuszno IG 1 borehole, 1729,7 m; Zechstein, PZ1 (Ca1) (x 1000).

Fig. 3. *Lueckisporites virkkiae* Potonié et Klaus NBb;

the Radwanów IG 1 borehole, 1644,5 m; Zechstein, PZ1 (Ca1) (x 1000).

Fig. 4. *Lueckisporites virkkiae* Potonié et Klaus NBC;

the Radwanów IG 1 borehole, 1660,0 m; Zechstein, PZ1 (Ca1) (x 1000).

Fig. 5. *Lueckisporites virkkiae* Potonié et Klaus NAc;

the Radwanów IG 1 borehole, 1729,7 m; Zechstein, PZ1 (Ca1) (x 1000).

Fig. 6. *Lueckisporites virkkiae* Potonié et Klaus NBa;

the Łopuszno IG 1 borehole, 1744,8 m; Zechstein, PZ1 (Ca1) (x 750).

Fig. 7. *Lueckisporites virkkiae* Potonié et Klaus f. with a "dark body" f. nov.; the Podgace IG 1 borehole, 97,7 m; Zechstein, PZ1 (T1r).

Fig. 8. *Lueckisporites virkkiae* Potonié et Klaus NE;
the Radwanów IG 1 borehole, 1644,5 m; Zechstein, PZ1 (Ca1) (x 1000).

Fig. 9. *Lueckisporites virkkiae* Potonié et Klaus NC (Guttulapollenites);
the Podgace IG 1 borehole, 124,7 m; Zechstein, PZ1 (Ca1) (x 750).

PLATE I




PLATE II




PLATE III




PLATE IV




PLATE V




PLATE VI




PLATE VII

